arXiv:0907.2407v1 [hep-th] 14 Jul 2009

D-independent representation of Conformal Field Theories in D dimensions via transformation to auxiliary Dual Resonance Models. Scalar amplitudes

G. Mack
II. Institut für Theoretische Physik, Universität Hamburg

July 14, 2009

Abstract

The Euklidean correlation functions and vacuum expectation values of products of n field operators $\phi^{k_i}(x_i)$ of some Lorentz spin l_i and dimension d_i are expressed through Mellin amplitudes $M_{k_n,\dots k_1}(\{\delta_{ij}\})$ which depend on complex dimensions $\delta_{ij} = \delta_{ji}$, $1 \le i < j \le n$ subject to linear constraints $\sum_j \delta_{ij} = d_i$. The constraints can be solved in terms of conserved momenta p_i whose squares are given by the field dimensions d_i, and related Mandelstam variables $s_{ij} = (p_i + p_j)^2$, viz. $\delta_{ij} = -p_i p_j$. The Mellin amplitudes furnish a universal representation of conformal field theories without explicit reference to D. The costumary principles of quantum field theory plus conformal invariance and operator product expansions (OPE) say that the Mellin amplitudes are amplitudes of dual resonance models with exact duality and a form of factorization which follows from OPE. It suffices to ensure these properties for all 4-point functions, together with positivity of all 2-point functions. A field with spin l and dimension d in the OPE's of scalar field products $\phi^{k_2}\phi^{k_1}$ and of $\phi^{k_3 *}\phi^{k_4 *}$ produces simple poles in the 4-point Mellin amplitude $M_{k_4 \dots k_1}$ at $s_{12} == d - l + 2n$, $n = 0, 1, 2....$ Their residues are polynomials of l-the order; they depend on D. The leading pole determines the satellites $n = 1, 2, 3, ...$

1 Introduction

There has recently been renewed interest in conformal field theories (CFT) in more than two space time dimensions D. Besides the perennial interest in the critical behavior of $D = 3$ dimensional models of statistical mechanics, including the Ising model, additional motivation comes from the AdS/CFT correspondence discovered by Maldacena [1, 9] which relates string theories to supersymmetric Yang Mills Theories in $D = 4$ dimensions. They are conformal field theories with the atypical property of admitting an expansion parameter, and are being studied vigorously [7]. CFT might also be useful in the description of unparticles [28].

Assuming Osterwalder Schrader positivity [57, 58], Euklidean Green functions, furnished e.g. by the correlation functions of a critical statistical mechanical system, can be analytically continued to Minkowski space and further to the ∞-sheeted cover $\mathcal{M}_D \simeq S^{D-1} \times \mathbf{R}$ of compactified Minkowski space [49]. This space admits a conformal invariant global causal ordering [48].

Conformal theories on compactified Minkowski space, whose fields have half-integral dimensions, are also being studied [56, 4, 55].

But here I am interested in theories with fields with anomalous dimensions.

The AdS/CFT correspondence also stimulated interest in relations between theories which live on space time manifolds of different dimension D (including relations between quantum field theories [65]). The known way how this can happen is through a holographic principle [68, 63]. I hope to pave the way for the study of different possibilities such as dimensional induction.

In conformal field theory, operator product expansions (OPE)[75, 76] have special properties. As noted first by Ferrara et.al [16, 17, 19], the contributions from derivatives of fields are determined by the others; a partial summation is therefore possible. The conformal OPE are partially summed forms of the Wilson OPE in which the contributions from all derivatives of a field are summed. Moreover, the conformal operator product expansions converge on the vacuum Ω.

This was proven in four dimensions [46] and also in two [39] and is generally true because it has a simple reason: It furnishes the orthogonal decomposition of the Hilbert space $\mathcal{H}$ of physical states in irreducible positive

energy representation spaces $\mathcal{H}^\chi$ of the conformal group [1]. The positive energy representations in 4 dimensions were determined in ref.[45].

A field ϕ with Lorentz spin l takes its values in a finite dimensional representation space V^l of the Lorentz group. Its components ϕ_α are obtained by expanding in a basis of V^l. I will mostly use vector notation.

In a conformal theory, the OPE take the form [46][2]

$$\phi^j(x)\phi^i(y)\Omega = \sum_k \sum_a g_{k,a}^{ji} \int d^D z \phi^k(z)\Omega \, \mathfrak{Q}^a(\chi_k, z; \chi_j, x, \chi_i, y) \tag{1}$$

The z-integration is over Minkowski space (or a conformal translate of it in the tube $\mathcal{M}_D$; the result is independent of the translation), and summation is over nonderivative fields.

$$\mathfrak{Q}^a(\chi_k, z; \chi_j, x, \chi_i, y) : V^{l_k} \mapsto V^{l_j} \otimes V^{l_i} \tag{2}$$

are *kinematically determined* generalized c-number functions (amputated three-point functions, see later). For scalar fields ϕ^j, ϕ^i there is only one of them, so that the index a and the sum over it can be dropped. The spin and dimension $\chi_k = [l_k, d_k]$ of all the fields and the coupling constants $g_{k,a}^{ij}$ are what has to be consistently determined to fix a theory, and in addition the normalizations $\mathbf{c}_k$ of some two point functions which have a physical meaning (i.e. for currents and the stress energy tensor).

There is the freedom of choosing a normalization factor in $\mathfrak{Q}$. The coupling constants g_k^{ji} depend on this choice. I will later adopt a particular choice of normalization; the kernel $\mathfrak{Q}(\chi...)$ with this normalization will be denoted by $\mathfrak{Q}^u(\tilde{\chi}...)$.

In any quantum field theory with operator product expansions which converge on the vacuum Ω, all the vacuum expectation values (n-point Wightman functions)

$$\begin{aligned} W_{i_n,...,i_1}(x_n, ..., x_1) &= \langle \Omega, \phi^{i_n}(x_n)...\phi^{i_1}(x_1)\Omega \rangle \\ &\in V^{l_{i_n}} \otimes ... \otimes V^{l_{i_1}} \end{aligned} \tag{3}$$

[1] It was conjectured in [47] that convergent operator product expansions exist also in the absence of conformal symmetry

[2] Throughout the paper, I use a special font, $\mathfrak{Q}$,$\mathfrak{V}$, $\mathfrak{M}$, to indicate *kinematically determined* quantities.

for arbitrary field operators $\phi^i(x)$ can be constructed from a consistent set of two- and three-point functions [47]. Consistency requires that all the 4-point functions obtained in this way satisfy locality. The four point functions possess a factorization property as a consequence of the OPE, see section 2 below, and they are positive if all 2-point functions are positive.

As shown in [47], any collection of 4-point functions with these properties defines a consistent quantum field theory. In CFT, the two and three-point functions of fields of given Lorentz spin and dimension are finite linear combinations of kinematically determined quantities [54, 53, 46]. The difficulty lies in satisfying locality.

All this was spelled out in detail long ago in the paper [47] on duality in quantum field theory, and it was also observed there that structures much reminiscent of dual resonance models emerge.

In particular, locality is a duality property. Here I derive more detailed results.

In conformal theories, the dynamical content is in functions of anharmonic ratios. Performing a Mellin transformation on these, one obtains a *Mellin representation* in which amplitudes $M^c_{i_n \dots i_1}(\{\delta_{ij}\})$ figure as coefficients. It represents the holomorphic function, whose boundary values are the Wightman functions, on their axiomatic domain of analyticity [67]. The *same* dual amplitude M^c enters the Mellin representation of the Euklidean Green functions, eqs.(47,48), the Wightman functions, eq.(50), and also the time ordered Green functions in Minkowski space, eq.(56). The latter two differ only through $i\epsilon$-prescriptions. The momentum space version of the Mellin representation of the time ordered Green function involves a generalized Feynman integral in the sense of Speer [66].

The Mellin amplitudes depend on complex dimensions $\delta_{ij} = \delta_{ji}$, $1 \leq i < j \leq n$ subject to linear constraints $\sum_i \delta_{ij} = d_j$. In the case $n = 4$, there are two independent variables. Using variables β_{ij} which differ from δ_{ij} by a shift, so that $\sum_i \beta_{ij} = 0$, β_{12} and $\gamma_{12} = \beta_{23} - \beta_{13}$ may be used.

I conjecture that the Mellin representation is also valid for massive field theories. The constraint $\sum_i \delta_{ij} = d_j$ on the variables δ_{ij} of integration is characteristic of conformal symmetry and will have to be relaxed in order to produce the subleading, i.e. less singular, contributions to the coefficients in the Wilson OPE. In contrast with scale invariant theories, these coefficients no longer scale like single powers of the distance in massive theories.

In this paper I show how the costumary principles of local relativistic quantum field theory plus conformal invariance and OPE, for 4-point func-

tions, translate into the statement that the Mellin amplitudes $M^c_{i_4\ldots i_1}$ have properties of dual resonance models.

Locality and some information from the OPE translate into exact duality, i.e. crossing symmetry and meromorphy in δ_{ij} with simple poles in single variables δ_{ij} [for n=4] whose residues are polynomials in the other independent variable(s). Duality is a D-independent property. CFT's in different dimensions D may have the same Mellin amplitude; for more on this see section 4.1. The precise relation between fields $\phi^k(x)$ of Lorentz spin l_k and dimension d_k which figure in the OPE and poles in the Mellin amplitudes depends on D and will be explained below. To every field there correspond a leading pole and satellites labeled by $n = 1, 2, ...$

I emphasize that the *meromorphy* of the dual amplitudes considered here is an *exact* property. In contrast to the dual resonance models of old, there is no "narrow resonance approximation" [74] involved, and no need to consider multiloop dual diagrams to satisfy unitarity [33]. The deeper reason for this is that the conformal two-point functions of fields ϕ^k usually have non-vanishing absorptive parts.

The interest in this result is in that there exist lots of examples of dual amplitudes; Veneziano's Beta-function Ansatz [73] was the first example.

Given duality, OPE translate into factorization properties of the residues of poles. Their precise form for scalar 4-point functions is described in section 9. Basically it is required that the residues of the leading poles factorize and the satellites are determined by them.

By the Osterwalder Schrader (OS) reconstruction theorem, both the spectrum condition and Wightman-positivity (unitarity) is satisfied if the Euklidean Green functions satisfy OS-positivity. Given duality and factorization, OS-positivity will hold if all the 2-point functions of the fields ϕ^k are OS-positive. This requires that $\chi_k = [l_k, d_k]$ label unitarizable representations of the confomal group, and it exorcizes $(-)$-signs in the factorization properties which signal ghosts.

Now I turn to the positions of the poles, for 4-point functions of scalar fields $\phi^{i_4}, ..., \phi^{i_1}$. I abbreviate $[l_{i_j}, d_{i_j}]$ by $[l_j, d_j]$.

If p_i are Lorentz vectors in any dimension, not necessarily equal to D, subject to the conservation law $\sum_{i=1}^n p_i = 0$, and with squares given by the dimensions of the external fields, $p_i^2 = d_i$, then the linear constraints $\sum_i \delta_{ij} = d_j$ can be solved in terms of Mandelstam variables

$$s_{ij} = (p_i + p_j)^2 \tag{4}$$

viz.

$$\delta_{ij} = -p_i p_j = -\frac{1}{2}(s_{ij} - d_i - d_j) \tag{5}$$

Consider a particular channel (21) $\mapsto$ (43) or, equivalently, the Wightman function $W_{i_4 \ldots i_1}$ with ordering of the fields as in eq.(3). Fields $\phi^k \neq \mathbf{1}$ with spin l_k and dimension d_k which appear in the OPE of both $\phi^{i_2}\phi^{i_1}$ and of $\phi^{i_3 *}\phi^{i_4 *}$ will lead to an integer spaced family of poles of $M_{i_4 \ldots i_1}$ in δ_{12}.

Stated in terms of the Mandelstam variables (5), poles in the channel (21) $\mapsto$ (43) are at positions

$$s_{12} = d_k - l_k + 2n, \qquad n = 0, 1, 2, \ldots \tag{6}$$

independent of the dimensions $d_1, \ldots, d_4$ of the external fields. This is the key result which permits the interpretation as dual resonance models, with twists $d_k - l_k$ substituting for mass squares. (For scalar fields $l_k = 0$.) To appreciate this, remember that in a 4-point scattering amplitude, poles in the Mandelstam variable $s_{12} = (p_1 + p_2)^2$ are determined by masses squared of resonances, *independent* of the masses of the particles which form the resonances.

To appreciate the role of field dimensions d_i, remember that a CFT with anomalous dimensions lives on $\mathcal{M}_D \simeq S^{D-1} \times \mathbf{R}$. There is a conformal Hamiltonian H which translates $\mathbf{R}$, and the radius of the compact space S^{D-1} furnishes a unit of length. The states $\phi^i(x)\Omega$ span an irreducible representation space labeled by $\chi_i = [l_i, d_i]$, and the eigenvalues of H in it are $d_i + n$, $n = 0, 1, 2, \ldots$ [49, 45]. Thus, d_i are energies in appropriate units.

What kinds of poles do we expect? It is expected that the fields which figure in the OPE, with Lorentz spin s and dimension d_s lie on trajectories $d_s = \alpha(s)$ involving fields with different s. These trajectories are what corresponds in quantum field theory to Regge-trajectories in S-matrix theory; they determine light cone singularities. It is expected that these trajectories are approximately, but not exactly, linear with slope 1. The evidence for this comes from the study of models which possess an expansion parameter. In ϕ^6-theory in $D = 6 + \epsilon$ dimensions [41], there is a fundamental real field of dimension $d = \frac{D}{2} - 1 + \Delta$, $\Delta = \frac{1}{18}\epsilon + \ldots$, and a trajectory of traceless symmetric tensor fields of even rank $s \geq 2$ with dimensions $d_s = D - 2 + s + \sigma_s$ whose anomalous part is given by the formula

$$\frac{1}{2}\sigma_s = \left[\frac{1}{18} - \frac{2}{3(s+2)(s+1)}\right]\epsilon + \ldots \tag{7}$$

so that $d_s - s \mapsto 2d$ as $s \mapsto \infty$. Callan and Gross showed that this property of the asymptotic behavior of d_s is always true in scalar field theory [8].

The member of the trajectory with $s = 0$ is absent. It is the shadow in the sense of Ferrara et al. [16] of the fundamental field. i.e. $d_0 = D - d$. The member $s = 2$ is the stress energy tensor; it has dimension $d_2 = D$.

The result (7) was obtained by conformal bootstrap. Expansions for anomalous dimensions in ϕ^4-theory in $4 - \epsilon$ dimensions and in N-vector models are also known [77, 71, 36, 72].

In the maximally supersymmetric Yang Mills theory in 4 dimensions, there is a trajectory whose anomalous part behaves as $\gamma \ln s$ for large s. This contribution is known as the cusp anomalous dimension; its coefficient γ was recently calculated [6].

Let us now examine what trajectories and satellites in the dual resonance model will correspond to the trajectories of fields ϕ^k. If $d_k = \alpha(l_k)$ then eq.(6) produces a trajectory of poles in the dual resonance model whose position in s_{12} is independent of the dimensions of the external fields

$$s_{12} = (\alpha(l_k) - l_k) + 2n, \tag{8}$$

$n = 0$, and satellites at $n = 1, 2, 3, ...$ We see that an exactly linear field trajectory of slope 1 would produce poles which fall on top of each other, i.e a linear trajectory of slope 0, plus satellites. In this paper, I am interested in fields with anomalous dimensions which lie on approximately linear field trajectories and I assume that there are at most finitely many poles in the Mellin amplitude which fall on top of each other. In a situation like (7), which is expected to occur in Ising-models, there will be limit points of poles. On the other hand, in the presence of a cusp anomalous part of the dimension, $\alpha(l_k) - l_k$ will go like $\gamma \ln l_k$ and there are no limit points. In either case, the CFT's correspond to a 0-slope limit of dual resonance models with only approximately linear trajectories.

It is a challenge to find a substitute for the operator formalism of old in dual resonance models [26] which ensures the appropriate form of factorization.

An alternative is to appeal to string theory. String theories were an outgrowth of dual resonance models [23, 31, 25, 32] and they are also 2-dimensional CFT's. Positivity in the CFT is needed to have a probability interpretation and would therefore be ensured by a consistent quantization of the string theory, i.e. a positive definite Hilbert space of its states, *if* the

state spaces can be identified. Identification of the states of the narrow resonances with states of a 2-dimensional CFT was an important tool in the old studies of dual resonance models. In the case of the Maldacena AdS/CFT -duality, the identification of the state spaces was established by group theoretical means, showing that the global symmetry and its representations are identical [9, 34]. The hope is, ultimately, that string theory will help to construct local relativistic quantum field theories. Exploiting relation (5), where p_i are Lorentz vectors of some dimension (determined by a string theory) with $p_i^2 = d_i$, one could try to get n-point Mellin amplitudes $M(\{\delta_{ij})$ from string field expectation values by an Ansatz like

$$\delta(\sum p_i) M(\{-p_i p_j\}) = \langle \int dV \; e^{i \sum_i p_{i\mu} X^\mu(\sigma_i, \tau_i)} \rangle \; . \tag{9}$$

The prototype of such a formula is in Veneziano's 1970 Erice lectures [74]. dV is an invariant volume element which integrates over $(n-3)$ of the arguments (σ_i, τ_i) of the string field X^μ. The time ordered Green functions of the D-dimensional CFT with scalar fields ϕ^i of dimension d_i are expressed in terms of M by eq.(56), viz.

$$\begin{aligned} &\langle \Omega, T\left\{\phi^n(x_n)...\phi^1(x_1)\right\}\Omega\rangle \\ &= (2\pi i)^{-n(n-3)/2} \int\int d\delta \; M(\{\delta_{ij}\}) \\ &\qquad \prod_{i>j} \Gamma(\delta_{ij})(-\frac{1}{2}x_{ij}^2 + i\epsilon)^{-\delta_{ij}}. \end{aligned}$$

Connected parts may have to be taken, see below.

To incorporate supersymmetry requires consideration of amplitudes with spin. This is postponed. The ultimate desire is to find an *explicit* duality transformation between quantum field theory at strong coupling and string theory at weak coupling.

There is a technical complication which affects the Mellin representation. It comes from the presence of disconnected parts and leads to an unexpected feature of the dual amplitudes. It is best stated in terms of the Euklidean Green functions $G_{i_n...i_1}(\mathbf{x}_n, ..., \mathbf{x}_1)$ which are the analytic continuation of the Wightman functions.

It is the full disconnected Greens function $G_{i_n...i_1}(\mathbf{x}_n, ..., \mathbf{x}_1)$ which satisfies the Osterwalder Schrader axioms, and to which operator product expansions apply.

In the dual resonance models of old, there are no disconnected parts, because they would correspond to processes $i, j \mapsto$ nothing in some channel, which is impossible on-shell.

The full disconnected Green functions decompose into a sum of disconnected parts and the connected Green functions $G^c_{i_n \ldots i_1}$. In case of the 4-point functions, the disconnected parts involve only two-point functions,

$$\begin{aligned} G_{i_4 \ldots i_1}(\mathbf{x}_n, ..., \mathbf{x}_1) &= G_{i_4 i_3}(\mathbf{x}_4, \mathbf{x}_3) G_{i_2 i_1}(\mathbf{x}_2, \mathbf{x}_1) + \\ & \quad G_{i_4 i_2}(\mathbf{x}_4, \mathbf{x}_2) G_{i_3 i_1}(\mathbf{x}_3, \mathbf{x}_1) + \\ & \quad G_{i_4 i_1}(\mathbf{x}_4, \mathbf{x}_1) G_{i_3 i_2}(\mathbf{x}_3, \mathbf{x}_2) + \\ & \quad G^c_{i_4 \ldots i_1}(\mathbf{x}_4, ..., \mathbf{x}_1) \ . \end{aligned} \tag{10}$$

There is one disconnected part for each channel (ji), e.g. (21) $\mapsto$ (43); it is given by the contribution of the unit operator $\mathbf{1}$ to the operator product $\phi^j \phi^i$ and vanishes if there is no such contribution.

For technical reasons, it is necessary to write a Mellin representation involving the dual amplitude M^c for the *connected* Green functions. Now consider the contribution from field ϕ^k in the operator product expansion of $\phi^{i_2} \phi^{i_1}$. It furnishes contributions to the asymptotic expansion of the full disconnected Green function $G_{i_4 \ldots i_1}(\mathbf{x}_4, ..., \mathbf{x}_1)$ when $\mathbf{x}_{12} = \mathbf{x}_1 - \mathbf{x}_2 \mapsto 0$. These contributions must match with contributions from the asymptotic expansion which obtains from the Mellin representation of the *connected* Green function, *plus* the disconnected parts. If $\phi^k = \mathbf{1}$, then the matching contribution is the disconnected part associated with channel (ji). Suppose $\phi^k \neq \mathbf{1}$. Then the matching contribution comes from the integer spaced family of poles (6) of M^c in δ_{12}.

However, if there are other non-vanishing disconnected parts than the afore mentioned one associated with channel (ji), then they must be cancelled by contributions from the Mellin representation of the connected Green function. For generic anomalous dimensions they would have the wrong degrees of homogeneity in x_{21} for contributions from OPE. Therefore the dual amplitude for the connected part must have kinematical poles in addition to the dynamical ones. They are fully determined in their positions and residues by the requirement that their contribution cancels the asymptotic expansion of disconnected parts (when they are nonzero and incompatible with OPE).

The Euklidean partial wave expansions suggest that these technical complications could be avoided by an illegitimate interchange of sums and integrations, which amounts to writing down a formal Mellin representation of

the full disconnected Green function. But I do not know how to do this in a legitimate way.

This paper deals with scalar amplitudes, mostly 4-point functions. n-point functions will be considered in section 3.3. The strategy of the paper is as follows. It would be difficult to find the polynomial residues of the dual amplitudes directly from the match between asymptotic expansions from OPE and from the Mellin representation. Therefore I will make a detour, using results established in the 70's on partial waves expansions of Euklidean Green functions (on the Euklidean conformal group $G^E = Spin(1, D+1)$), and their relation to OPE in Minkowski space [42, 43, 12, 11, 14]. As a check of the result, I will verify that the match between asymptotic expansions obtains for the special case of the contribution of a scalar field to the OPE and the associated family of poles in the Mellin amplitude.

The dynamical information on the scalar four point function $G_{i_4 i_3 i_2 i_1}$ within the Euklidean partial wave expansion in channel $(21) \mapsto (43)$ resides in a complex function $g^{(43)(21)}(\chi)$ of $\chi = [l, \delta]$, where l is again a (traceless symmetric tensor) representation of the (Euklidean) Lorentz group, and δ is a complex dimension. It has the symmetry property

$$g^{(43)(21)}(\chi) = g^{(43)(21)}(\tilde{\chi})$$

where $\tilde{\chi} = [\bar{l}, D - \delta]$, with $\bar{l}$ the dual representation to l. Validity of OPE expansions requires that $g^{(43)(21)}(\chi)$ is a meromorphic function of δ and there is a bijective correspondence between (pairs of) poles of $g^{(43)(21)}(\chi)$ and fields ϕ^k in the OPE as follows. A field with Lorentz spin and dimension $\chi_k = [l_k, d_k]$ and which contributes to the OPE (1) of $\phi^{i_2}\phi^{i_1}$, with coupling constant g_k^{21} *and* of $\phi^{i_3 *}\phi^{i_4 *}$, with coupling constant $\bar{g}_k^{43}$ requires a pole of $g^{(43)(21)}([l_k, \delta])$ at $\delta = d_k$ with residue

$$\bar{g}_k^{43} g_k^{21} \mathbf{c}_k \tag{11}$$

times a kinematical factor. $\mathbf{c}_k$ is given by the normalization of the 2-point function of ϕ^k. I will first derive the Mellin representation of individual Euklidean conformal partial waves, using an integration formula of Symanziks [70], and then deduce the meromorphy properties of the dual amplitude, including its residues, from the meromorphy properties of $g^{(43)(21)}(\chi)$. The factorization properties of the dual amplitude come from the factorized expression (11) of the residue of $g^{(43)(21)}([l_k, \delta])$.

2 Positivity

2.1 Fields and positive energy representations

I consider collections of nonderivative field operators $\phi^i(x)$, including the unit operator $\mathbf{1}$, which are closed under OPE. The conformal transformation law of ϕ^i is specified by

$$\chi_i = [l_i, d_i] \tag{12}$$

where d_i are positive real, called the dimension of the field ϕ^i, and l_i specifies an irreducible finite dimensional representation of the Lorentz group M by operators $D^{l_i}(m)$ in a vector space V^{l_i}. I write $l = 0$ for the trivial 1-dimensional representation of M. Scalar fields have $l = 0$. For the unit operator, $l = 0$, $d = 0$. Physically allowed values of χ are in bijective correspondence with unitary irreducible positive energy representations of the conformal group G in D dimensions. They depend on D. Fields $\phi^i_\alpha(x)$ with spinor indices α are obtained by expanding the vector $\phi^i(x) \in V^{l_i}$ in a multispinor basis of V^{l_i}.

For the sake of transparency, I will write multispinor indices α of fields $\phi^i_\alpha(x)$ in this section.

When anomalous (i.e. not half integral) field dimensions are admitted, CFT lives on the ∞-sheeted simply connected universal covering $\mathcal{M}_D$ of conformally compactified Minkowski space, and G is the infinite sheeted universal covering of $SO(2, D)$ [49]. The Wightman functions are determined by their values on Minkowski space, but admit unique analytic continuation to $\mathcal{M}_D$ and appropriately defined field operators may be regarded to live on $\mathcal{M}_D$. State vectors of the special form $\phi^k_\alpha(z)\Omega$ span irreducible representation spaces $\mathcal{H}^k$ of G. These state vectors are periodic on $\mathcal{M}_D$ so that they can be regarded as sections of vector bundles over compactified Minkowski space. (In even dimensions this follows from Schur's lemma, applied to the center of G). In the following, all integrations $\int d^D z$ are over Minkowski space.

2.2 Positivity from Operator Product Expansions

I demand the standard properties of Wightman functions [67] - spectrum condition, positivity and locality plus conformal invariance, and validity of operator product expansions (OPE) (1) which converge on the vacuum Ω.

Let us analyze positivity. Consider finite sequences f of test functions on

Minkowski space,

$$f^{\emptyset}, f^{i_1}_{\alpha_1}(x_1), \ldots, f^{i_1 \ldots i_N}_{\alpha_1 \ldots \alpha_N}(x_1, \ldots, x_N)$$

where $f^{\emptyset}$ is a constant. For given sequence f, consider the state vector

$$\begin{aligned} \Psi(f) &= f^{\emptyset}\Omega + \sum_{n=1}^{N} \int d^D x_n ... \int d^D x_1 f^{i_1 \ldots i_n}_{\alpha_1 \ldots \alpha_n}(x_1 ... x_n) \\ & \qquad \phi^{i_n}_{\alpha_n}(x_n) \ldots \phi^{i_1}_{\alpha_1}(x_1)\Omega \ . \end{aligned} \tag{13}$$

Summation over all repeated indices $\alpha_1, ..., i_n$ is understood. $\Psi(f)$ is an element of the Hilbert space of physical states, and must therefore have positive semidefinite norm,

$$\langle \Psi(f), \Psi(f) \rangle \geq 0 \tag{14}$$

Inserting the definition (13) one obtains an inequality on Wightman functions [67] which is known as Wightman positivity. It is related to unitarity in the language of time ordered Green functions.

The analysis of Wightman positivity is much simplified by validity of OPE. Consider first sequences with $N = 2$. Inserting in definition (13) the OPE (1) for $\phi^{i_2}_{\alpha_2}(x_2)\phi^{i_1}_{\alpha_1}(x_1)\Omega$ we obtain

$$\Psi(f) = \sum_{k}{}' \int d^D z h^k_{\alpha}(z) \phi^k_{\alpha}(z)\Omega + h^{\emptyset}\Omega \tag{15}$$

Summation $\sum'$ is over all non-derivative fields *other than* the unit operator. Explicitly, the functions $h^k_{\alpha}(z)$ are given by

$$\begin{aligned} h^k_{\alpha}(z) &= \sum_{a}{}' g^{i_2 i_1}_{k,a} \int \int d^D x_2 d^D x_1 f^{i_1 i_2}_{\alpha_1 \alpha_2}(x_1, x_2) \\ & \qquad \mathfrak{Q}^a_{\alpha \alpha_2 \alpha_1}(\chi_k, z; \chi_{i_2}, x_2, \chi_{i_1}, x_1) + f^k_{\alpha}(z) \\ h^{\emptyset} &= g^{i_2 i_1}_0 \int \int d^D x_2 d^D x_1 f^{i_1 i_2}_{\alpha_1 \alpha_2}(x_1, x_2) \\ & \qquad \mathfrak{Q}_{\emptyset \alpha_2 \alpha_1}(0, z; \chi_{i_2}, x_2, \chi_{i_1}, x_1) + f^{\emptyset} \ . \end{aligned}$$

Here I write $\chi = 0$ for the trivial 1-dimensional representation and $\alpha = \emptyset$ when there is no index. The kernel $\mathfrak{Q}$ in the expression for $h^{\emptyset}$ is actually a

two-point function independent of z. There is only one such, therefore there is no index a to sum over in the formula for $h^{\emptyset}$.

Inserting eq.(15), and using that in a conformal theory $\langle \Omega, \phi^k_\alpha(z)\Omega\rangle = 0$ for all fields ϕ^k other than the unit operator, we obtain

$$\begin{aligned} \langle \Psi(f), \Psi(f)\rangle &= \sum_k{}' \sum_{k'}{}' \int\int d^D z d^D z' \qquad (16) \\ &\quad \bar{h}^k_\alpha(z)\langle \Omega, \phi^{k*}_\alpha(z)\phi^{k'}_{\alpha'}(z')\Omega\rangle h^{k'}_{\alpha'}(z') \\ &\quad + \bar{h}^{\emptyset} h^{\emptyset}\langle \Omega, \Omega\rangle \qquad (17) \end{aligned}$$

This is positive semidefinite *provided* the two-point functions of all the fields are positive.

Using the OPE repeatedly as described in ref. [47] it is seen that eq.(15) generalizes to arbitrary finite sequences f, and the same is therefore also true of the result (16).

Conformal invariant two point functions are determined by conformal symmetry up to an overall factor. Taking linear combinations of fields if necessary,

$$\langle \Omega, \phi^{k*}_\alpha(z)\phi^{k'}_{\alpha'}(z')\Omega\rangle = \mathbf{c}_k \delta_{kk'} \Delta^{\chi_k}_{\alpha\alpha'}(z, z') \qquad (18)$$

where $\Delta^{\chi_k}(z, z')$ is kinematically determined. If the field transformation law is given by $\chi_k = [l_k, d_k]$ which determines a unitarizable representations, then Δ^{χ_k} is positive and the two point functions are either positive or negative semidefinite, depending on whether the overall factor $\mathbf{c}_k$ is positive or negative. Fields whose two-point function is negative are called *ghosts*.

In conclusion, once validity of OPE is assured and if the fields ϕ^k have spin and dimension $\chi_k = [l_k, d_k]$ which correspond to unitary representations of G, positivity is assured if none of the fields is a ghost.

The requirement that χ_k should label unitary positive energy representations of G will be called the requirement of *no tachyons*. Actually it also comes out of the requirement of positivity. But it is convenient to divide positivity into two requirements: Absence of tachyons which gives D-dependent restrictions on the allowed positions of poles of Mellin amplitudes, and absence of ghosts which could be inherited from a 2-dimensional theory.

2.3 Factorization properties from Operator Product Expansions

Consider two pairs of fields, labeled by j, i and l, m. Assume that fields are labeled in such a way that $\phi^{k*} = \phi^{\bar{k}}$.

Inserting the OPE (1) for $\phi^j\phi^i\Omega$ and for $\phi^l\phi^m\Omega$, and using the formula (18) for the conformal invariant two-point functions, the OPE for the four-point Wightman functions takes the form

$$\begin{aligned} W_{\bar{m},\bar{l},j,i}(x_4,...,x_1) &= \langle\Omega, \phi^{\bar{m}}(x_4)\phi^{\bar{l}}(x_3)\phi^j(x_2)\phi^i(x_1)\Omega\rangle \\ &= \sum_k \sum_{a,b} \mathbf{c}_k \bar{g}^{lm}_{k,b} g^{ji}_{k,a} \mathfrak{W}^{ba}_{\chi_k,\bar{m},\bar{l},j,i}(x_4,...,x_1) \end{aligned} \tag{19}$$

etc., where $\mathfrak{W}$ are kinematically determined quantities,

$$\begin{aligned} \mathfrak{W}^{ba}_{\chi_k,\bar{m},\bar{l},j,i}(x_4,...,x_1) = \int\int dz\ dz' \\ \bar{\mathfrak{Q}}^b(\chi_k,z;\chi_l,x_3,\chi_m,x_4)\Delta^{\chi_k}(z,z')\mathfrak{Q}^a(\chi_k,z';\chi_j,x_2,\chi_i,x_1) \end{aligned} \tag{20}$$

Suppose that a particular field ϕ^k contributes to both $\phi^j\phi^i\Omega$ and $\phi^l\phi^m\Omega$. Then it will contribute to all of the following Wightman functions and its contributions are related as follows.

Assume for transparency that there is only one invariant 3-point function so that the label a on $\mathfrak{Q}^a$ in eq.(1) is redundant. Then the contribution of the field ϕ^k to the 4-point Wightman functions is

$$\begin{aligned} W_{\bar{m},\bar{l},j,i}(x_4,...,x_1) &= \mathbf{c}_k \bar{g}^{lm}_k g^{ji}_k \mathfrak{W}_{\chi_k,\bar{m},\bar{l},j,i}(x_4,...,x_1) \\ &\quad + \ldots, \\ W_{\bar{i},\bar{j},j,i}(x_4,...,x_1) &= \mathbf{c}_k \bar{g}^{ji}_k g^{ji}_k \mathfrak{W}_{\chi_k,\bar{i},\bar{j},j,i}(x_4,...,x_1) \\ &\quad + \ldots, \\ W_{\bar{m},\bar{l},l,m}(x_4,...,x_1) &= \mathbf{c}_k \bar{g}^{lm}_k g^{lm}_k \mathfrak{W}_{\chi_k,\bar{m},\bar{l},l,m}(x_4,...,x_1) \\ &\quad + \ldots. \end{aligned}$$

The real constants $\mathbf{c}_k$ are determined by the two-point functions (18). They are positive in a theory without ghosts.

We see that the coefficients of the kinematically determined factors $\mathfrak{W}$ factorize in a product of coupling constants in a way which is familiar from dual resonance models. These factorization properties are at the origin of the

factorization properties of the residues of the poles in the Mellin amplitude, and they assure positivity under the above mentioned conditions.

For future use we note that the kinematically determined factors $\mathfrak{W}$ can be written in another way.

Consider three-point functions. There is a finite number of kinematically determined conformal invariant 3-point functions $\mathfrak{V}^a$, so that

$$\begin{aligned} &\langle\Omega, \phi^k(x_3)\phi^j(x_2)\phi^i(x_1)\Omega\rangle \\ &= \sum_a \tilde{g}^{ji}_{k,a}\mathfrak{V}^a(\chi_k, x_3, \chi_j, x_2, \chi_i, x_1) \end{aligned} \tag{21}$$

with dynamically determined coupling constants $\tilde{g}^{ji}_{k,a}$ which will be related to $g^{ji}_{k,a}$ below.

Inserting the OPE in the 3-point function and using (18) gives

$$\begin{aligned} \langle\Omega, \phi^{\bar{k}}(x_3)\phi^j(x_2)\phi^i(x_1)\Omega\rangle &= \mathbf{c}_k \sum_a \int dz \Delta^{\chi_k}(x_3, z) \\ &\quad g^{ji}_{k,a}\mathfrak{Omega}^a(\chi_k, z; \chi_j, x_2, \chi_i, x_1) \end{aligned} \tag{22}$$

Exploiting the freedom of taking linear combinations of the kinematically determined quantities we may demand the following *amputation identity*

$$\begin{aligned} &\mathfrak{V}^a(\chi_{\bar{k}}, x_3, \chi_j, x_2, \chi_i, x_1) = \\ &\int dz.\Delta^{\chi_k}(x_3, z)\mathfrak{Omega}^a(\chi_k, z; \chi_j, x_2, \chi_i, x_1) \end{aligned} \tag{23}$$

. and

$$\tilde{g}^{ji}_{k,a} = \mathbf{c}_k g^{ji}_{k,a} \tag{24}$$

This shows that $\mathfrak{Omega}$ is an amputated Wightman 3-point function. It is uniquely determined by this and by the requirement that its partial Fourier transform in z is a holomorphic function of p.

Inserting eq.(22 simplifies the defining formula (20) of $\mathfrak{W}$. The resulting formula will be used in section 11 for four-point functions of real scalar fields; the labels a, b are redundant in this case, and $\bar{l} = l$ etc..

$$\begin{aligned} &\mathfrak{W}^{be}_{\bar{m},\bar{l},j,i}(x_4, ..., x_1) = \\ &\int dz.\bar{\mathfrak{Omega}}^b(\chi_k, z; \chi_l, x_3, \chi_m, x_4)\mathfrak{V}^a(\chi_k, z, \chi_j, x_2, \chi_i, x_1) \\ = \quad &\mathfrak{Omega}^b(\chi_k, p; \chi_m, x_4, \chi_l, x_3)\mathfrak{V}^a(\chi_k, z, \chi_j, x_2, \chi_i, x_1)|_{z=0} \ . \end{aligned} \tag{25}$$

. In the last formula, $\mathfrak{Q}(\chi_k, p; ...)$ is the partial Fourier transform of $\mathfrak{Q}$ with respect to the variable z; it has the reality property (87). p is to act as a differential operator with respect to z, $p = -i\nabla_z$. As noted before, indices a, b are redundant when scalar four-point amplitudes are expanded.

3 Universal Mellin representation

3.1 Euklidean Green functions

Wightman functions admit analytic continuation to Euklidean points x, defining Euklidean Green functions

$$G_{i_n...i_1}(\mathbf{x}_n, ..., \mathbf{x}_1) \in V^{l_n} \otimes .. \otimes V^{l_1}. \tag{26}$$

Euklidean points x have imaginary time components x^0 and real space components x^i, $i = 1, ..., D-1$. They can also be specified by real coordinates $\mathbf{x} = (x^1, ..., x^D)$, $x^D = ix^0$. *Whenever x and $\mathbf{x}$ will appear in the same formula, they are understood to be related in this way.* In particular,

$$-x^2 = \mathbf{x}^2 > 0$$

in Euklidean space.

By Weyls unitary trick, the finite dimensional representation spaces V^l of the Lorentz group are identified with representation spaces of its Euklidean brother $Spin(D)$. The Euklidean Green functions so defined include disconnected parts, and satisfy Oterwalder Schrader positivity as a substitute for spectrum condition and positivity of Wightman functions. Locality becomes (crossing) symmetry,

$$G_{i_{\pi n}...i_{\pi 1}}(\mathbf{x}_{\pi n}, ..., \mathbf{x}_{\pi 1}) = \pm\, \hat{\pi}\; G_{i_n...i_1}(\mathbf{x}_n, ..., x_1) \tag{27}$$

for arbitrary permutations π of $1...n$. $\pm$ depends on how many Fermi fields are interchanged, and

$$\hat{\pi} : V^{l_n} \otimes .. \otimes V^{l_1} \mapsto V^{l_{\pi n}} \otimes .. \otimes V^{l_{\pi 1}}$$

is the map which permutes multispinor indices. It is redundant for scalar Green functions.

The Euklidean Green functions decompose into connected Greens functions plus possibly disconnected parts as in eq.(10). The above symmetry property is shared by the connected Greens functions.

The Eukidean Green functions are invariant under the Euklidean conformal group $Spin(D+1,1)$ which is isomorphic to the Lorentz group in $D+2$ dimensions.

3.2 Mellin representation of scalar 4-point functions

Let us first consider Euklidean 4-point Green functions of four scalar fields of dimensions $d_1, ..., d_4$ including in particular the Greens function $G_{0000}(\mathbf{x}_4, ..., \mathbf{x}_1) \in \mathbf{C}$ of a single scalar field ϕ^0 of dimension d_0. For the latter, the symmetry property simplifies to

$$G_{0000}(\mathbf{x}_{\pi 4}, ..., \mathbf{x}_{\pi 1}) = G_{0000}(\mathbf{x}_4, ..., \mathbf{x}_1). \tag{28}$$

The arbitrariness in a scalar conformal invariant 4-point functions resides in an arbitrary function $F(\omega)$ of anharmonic ratios. They are defined as follows, with $x_{ij} = x_i - x_j$.

$$\omega_1 = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}, \quad \omega_2 = \frac{x_{14}^2 x_{23}^2}{x_{12}^2 x_{34}^2}, \quad \omega_3 = \frac{x_{13}^2 x_{24}^2}{x_{14}^2 x_{23}^2}, \tag{29}$$

and satisfy

$$\omega_1 \omega_2 \omega_3 = 1 \ . \tag{30}$$

For non-coinciding Euklidean points,

$$x_{ij}^2 = -|\mathbf{x}_{ij}|^2 < 0$$

and all the anharmonic ratios are positive. We write $\omega = (\omega_1, \omega_2, \omega_3)$ for triples of anharmonic ratios subject to (30). Permutations π of arguments $\mathbf{x}_1 ... \mathbf{x}_4$ act on such triples ω as follows. Consider transpositions (ij) of two arguments i and j.

$$\begin{aligned}
&(12) \text{ or } (34): \quad \omega_1 \mapsto \omega_2^{-1}, \quad \omega_2 \mapsto \omega_1^{-1}, \quad \omega_3 \mapsto \omega_3^{-1}, \\
&(13) \text{ or } (24): \quad \omega_1 \mapsto \omega_3^{-1}, \quad \omega_2 \mapsto \omega_2^{-1}, \quad \omega_3 \mapsto \omega_1^{-1}, \\
&(14) \text{ or } (23): \quad \omega_1 \mapsto \omega_1^{-1}, \quad \omega_2 \mapsto \omega_3^{-1}, \quad \omega_3 \mapsto \omega_2^{-1},
\end{aligned}$$

Note that some even permutations of x_i act trivially on $(\omega_1, \omega_2, \omega_3)$,

$$\pi\omega = \omega \text{ for } \pi = (12)(34) \text{ or } (13)(24) \text{ or } (14)(23) \ . \tag{31}$$

The other even permutations of x_i act as cyclic permutations of $(\omega_1, \omega_2, \omega_3)$.

Consider complex exponents $\delta_{ij} = \delta_{ji}$ attached to unordered pairs (ij) of distinct $i, j = 1...4$ subject to the constraint

$$\sum_i \delta_{ij} = d_j \tag{32}$$

There is a unique real solution $\{\delta^0_{ij} = \delta^0_{ij}(d_1, ..., d_n)\}$ of this system of equations which satisfies the condition

$$\delta^0_{\pi i \pi j}(d_{\pi 1}, ..., d_{\pi n}) = \delta^0_{ij}(d_1, ..., d_n) \tag{33}$$

For $n = 4$ it is given by

$$\delta^0_{12} = \frac{1}{6}(2d_1 + 2d_2 - d_3 - d_4) \text{ etc.} \tag{34}$$

Given the dimensions $d = \{d_i\}$ of the four scalar fields, $i = 1...4$ define the *fundamental scalar 4-point function*

$$f(x_1, ..., x_4|d) = \prod_{ij}(-\frac{1}{2}x^2_{ij})^{-\delta^0_{ij}} \tag{35}$$

The most general conformal invariant Euklidean scalar 4-point Green function has the form

$$G_{i_4...i_1}(\mathbf{x}_1, ..., \mathbf{x}_4) = f(x_1, ..., x_4|d)F_{i_4...i_1}(\omega) \tag{36}$$

and locality reads

$$F_{i_{\pi 4}...i_{\pi 1}}(\pi\omega) = F_{i_4...i_1}(\omega) \tag{37}$$

In particular, $F_{0000}(\pi\omega) = F_{0000}(\omega)$, and these relations depend neither on D nor on the dimensions d_i of the scalar fields.

We also see that locality condition (28) is automatically fulfilled for those permutations π, given by eq.(31), for which $\pi\omega = \omega$.

In general, the full disconnected Green function is the sum of disconnected parts and a connected Green function G^c, as in eq.(10). This translates into sums of terms of the form (36) in the obvious way. For the amplitude G_{0000} of four equal fields of dimension d_0 with two-point function

$$G_{00}(\mathbf{x}_2, \mathbf{x}_1) = N_{d_0}(-\frac{1}{2}x^2_{12})^{-d_0},$$

the decomposition reads

$$F_{0000}(\omega) = N_{d_0}^2 \left(\left(\frac{\omega_2}{\omega_1}\right)^{d_0/3} + \left(\frac{\omega_1}{\omega_3}\right)^{d_0/3} + \left(\frac{\omega_3}{\omega_2}\right)^{d_0/3} \right) + F_{0000}^c(\omega) \ . \tag{38}$$

Note that the sum of disconnected parts fulfills locality constraint (37) as it must be. In a generalized free field theory, $F^c_{i_4 \dots i_1} = 0$.

Now we subject $F^c(\omega)$ to Mellin transformation. Given the dimensions of fields, OPE yield information about the strength of the singularities of the Euklidean Green functions at coinciding arguments, and this translates into information on the strength of the singularities of $F(\omega)$ when some $\omega_a \mapsto 0$ or ∞. If these singularities are not too strong, and if some further boundedness properties hold, a Mellin expansion exists with paths that can be taken along the imaginary axis [3]. It is convenient to write the expansion in the following form.

The general solution of eq.(32) is of the form

$$\delta_{ij} = \delta^0_{ij} + \beta_{ij} \tag{39}$$

where $\beta = \{\beta_{ij}\}$ is a solution of the homogeneous equation, whence

$$\beta_{34} = \beta_{12}, \quad \beta_{24} = \beta_{13}, \quad \beta_{14} = \beta_{23}, \tag{40}$$

$$\beta_{12} + \beta_{13} + \beta_{23} = 0 \ . \tag{41}$$

It depends on two independent parameters, e.g. β_{12} and β_{13}. Abbreviating $(i_4 ... i_1) = a$ to specify a quadruple of scalar fields, the Mellin expansion reads

$$F^c_a(\omega) = (2\pi i)^{-2} \int_{-i\infty}^{i\infty} \int_{-i\infty}^{i\infty} d^2\beta \ \hat{M}^c_a(\beta) \omega_1^{-s_1} \omega_2^{-s_2} \omega_3^{-s_3}, \tag{42}$$

where s_i are determined by β up to a common additive constant, which is irrelevant by (30), and

$$s_2 - s_1 = -\beta_{12}, \quad s_3 - s_2 = -\beta_{23}, \quad s_1 - s_3 = -\beta_{13} \tag{43}$$

[3] It is known that an expansion of this form exists order by order in a skeleton graph expansion, for Greens functions of fundamental fields. It can be derived using Symanzik's integration formula (143)

$d^2\beta$ is integration over two independent variables, e.g. $d^2\beta = d\beta_{12}d\beta_{13}$.

$\hat{M}^c_a$ will be called the *reduced Mellin amplitude*. There is an obvious action of permutations π of the four space time points on β, viz.

$$(\pi\beta)_{ij} = \beta_{\pi^{-1}i\pi^{-1}j} \tag{44}$$

and locality reads

$$\hat{M}_{\pi a}(\pi\beta) = \hat{M}_a(\beta) \tag{45}$$

There is neither an explicit reference to D nor to the dimensions of the scalar fields.

Defining, for $\delta_{ij} = \delta^0_{ij} + \beta_{ij}$ the *Mellin amplitude* M^c by

$$M^c_a(\{\delta_{ij}\}) = \hat{M}^c_a(\{\beta_{ij}\}) \prod_{ij} \Gamma(\delta_{ij})^{-1}, \tag{46}$$

the connected scalar 4-point function can be written as

$$\begin{aligned} G^c_a(\mathbf{x}_4, ..., \mathbf{x}_1) &= (2\pi i)^{-2} \int\int d\delta \qquad M^c_a(\{\delta_{ij}\}) \\ &\qquad \prod_{ij} \Gamma(\delta_{ij}) (-\frac{1}{2} x^2_{ij})^{-\delta_{ij}}, \end{aligned} \tag{47}$$

with $a = (i_4...i_1)$, $d\delta = d^2\beta = d\beta_{12}d\beta_{13}$ and integration over imaginary β_{12}, β_{13}. In contrast with the reduced Mellin amplitude, M^c depends on the dimensions of the scalar fields.

The explicit Γ-functions decrease exponentially in the imaginary directions. They are included in order that the growth conditions on the Mellin amplitude, which ensure the axiomatic analyticity in coordinate space, take a simple form; cf. section 3.4.

It will be shown in section 8 that OPE in some channel, let us say (21) $\mapsto$ (43), determine the poles of M^c in δ_{12} (and δ_{34}) and their residues. But there are also zeroes at non-positive integers δ_{12} and δ_{34} so that the reduced Mellin amplitude $\hat{M}^c$, which is related to M^c by eq.(46), does not have extra poles at non-positive integers δ_{12} and δ_{34}. The result suggests to regard the factor M^c in the Mellin representation (47) as an operator which creates the integrand of (47) from the raw material $\prod \Gamma(\delta_{ij})(-\frac{1}{2}x^2_{ij})^{-\delta_{ij}}$ by shifting poles and modifying residues.

3.3 Scalar n-point functions for $n > 4$

The generalization of the Mellin representation to $n > 4$ is straightforward. Given the field dimensions d_i, $i = 1...n$, consider again solutions $\delta_{ij} = \delta_{ji}$, $j \neq i$ of the system of equations

$$\sum_i \delta_{ij} = d_j \qquad j = 1 \dots n$$

The solution space is $\frac{1}{2}n(n-3)$ dimensional [70]. There is a unique solution δ^0_{ij} of this system of equations with the property (33), and the general solution is $\delta_{ij} = \delta^0_{ij} + \beta_{ij}$, where β_{ij} is the general solution of the homogeneous equation; $\frac{1}{2}n(n-3)$ of these variables are independent. We may choose β_{ij} with $2 \leq i < j < n$, excepting δ_{23} as the independent ones [70]. The Mellin representation of the connected Green function reads

$$\begin{aligned} G^c_a(\mathbf{x}_n, ..., \mathbf{x}_1) &= (2\pi i)^{-n(n-3)/2} \int\int d\delta \; M^c_a(\{\delta_{ij}\}) \\ &\prod_{ij} \Gamma(\delta_{ij})(-\frac{1}{2}x^2_{ij})^{-\delta_{ij}}, \end{aligned} \tag{48}$$

with $a = (i_n, ..., i_1)$ and $d\delta = d^{n(n-3)/2}\beta$. A reduced Mellin amplitude $\hat{M}^c(\beta)$ can be defined by eq. (46).

The n-point functions and their Mellin amplitudes can be constructed from two- and three-point functions, and these in turn are furnished by a collection of factorizing local four-point functions. The meromorphy and factorization properties of the n-point Mellin amplitudes follow from this. The coordinate space integrations which are involved in constructing n-point functions from lower ones can be converted into operations on the Mellin amplitudes by using Symanzik's integration formula with spin.

Abbreviating $\delta = \{\delta_{ij}\}$, retaining $a = (i_n, ..., i_1)$, and defining $(\pi\delta)_{ij} = \delta_{\pi^{-1}i\pi^{-1}j}$, locality reads

$$M^c_{\pi a}(\pi\delta) = M^c_a(\delta) \tag{49}$$

3.4 Spectrum condition

Given suitable growth conditions on M^c_a, the right hand sides of the Mellin representations (47,48) also defines the analytic continuation of the Euklidean Green function throughout the axiomatic analyticity domain, known

as the permuted extended tube, and also its boundary values, the Wightman functions, when the appropriate $i\epsilon$-prescriptions are understood which are appropriate for a particular ordering of the fields.

The $i\epsilon$-prescriptions assure validity of the spectrum condition, and follow from it.

Consider the Wightman function which is defined by a particular $i\epsilon$-prescription, and its Fourier transform

$$\begin{aligned}
&\langle\Omega, \phi^{i_n}(x_n)...\phi^{i_1}(x_1)\Omega\rangle = \\
&(2\pi)^{-nD}\int ...\int\left(\prod_{i=1}^{n} dp_i e^{-ip_i x_i}\right) W_a(p_n,...,p_1) \\
&= (2\pi i)^{-n(n-3)/2}\int\int d\delta\ M_a^c(\{\delta_{ij}\}) \\
&\qquad \prod_{i>j}\Gamma(\delta_{ij})(-\frac{1}{2}x_{ij}^2 + i\epsilon x_{ij}^0)^{-\delta_{ij}}, \qquad (50)
\end{aligned}$$

with the abbreviation $a = (i_n, ..., i_1)$.

The spectrum condition says that all physical states must have D-momentum p in the closed forward light-cone V_+. This is equivalent [67] to the requirement that $W(p_n, ..., p_1)$ vanishes unless

$$q_k \equiv -\sum_{i=1}^{k} p_i \in V_+ \qquad (51)$$

for all $k = 1...n-1$. Now insert the integral representation of the factors

$$\Gamma(\delta)(-\frac{1}{2}x^2 + i\epsilon x^0)^{-\delta} \;=\; \rho_D(\delta)\int_{V_+} dp e^{-ipx}\left(\frac{p^2}{2}\right)^{-h+\delta}\ , \qquad (52)$$

with

$$\rho_D(\delta) = 2^{-2\delta+1}\pi^{-h+1}\Gamma(\delta)[\Gamma(2h-\delta)\Gamma(\delta-h+1)]^{-1}\ ,$$

$h = \frac{1}{2}D$. Define variables of integration p_{ij} for $1 \le j < i \le n$, one for each unordered pair (ij), and set $p_{ij} = -p_{ji}$ for $i < j$, and $p_{ii} = 0$. Inserting the integral representation (51) for each x-dependent factor in expression (50), one can read off the Fourier transform

$$\begin{aligned}
&(2\pi)^{nD}W(p_n,...,p_1) = \qquad (53) \\
&\int_{V_+} ...\int_{V_+}\prod_{i>j} dp_{ij}\prod_k \delta(p_k - \sum_{m=1}^{n} p_{km})\tilde{M}^c(\{p_{ij}\})
\end{aligned}$$

with

$$\tilde{M}(\{p_{ij}\}) \;=\; (2\pi i)^{-n(n-3)/2} \int\int d\delta \; M^c(\{\delta_{ij}\}) \prod_{i>j} \rho_D(\delta_{ij}) \left(\frac{p_{ij}^2}{2}\right)^{-h+\delta_{ij}} \tag{54}$$

From the δ-functions it follows that

$$-\sum_{i=1}^{k} p_i = \sum_{j=k+1}^{n} \sum_{i=1}^{k} p_{ji} \;\in\; V_+$$

throughout the domain of integration. Therefore the spectrum condition is fulfilled.

Conversely, it follows [67] from the spectrum condition that the Wightman function, considered as a function of differences of coordinates, is holomorphic in the half planes $Im\ x_{ij}^0 < 0$, for $i > j$. The $i\epsilon$ prescription shifts infinitesimally into this half plane since $-\frac{1}{2}(x^0 - i\epsilon)^2 = -\frac{1}{2}(x^0)^2 + i\epsilon x^0$.

Concerning the growth conditions, we do not need to worry about them once we have Osterwalder Schrader positivity as a consequence of OPE, see section 5.6. Since the other Euklidean axioms are fulfilled by construction, the Osterwalder Schrader reconstruction theorem can then be applied [57, 58]. It will guarantee validity of the spectrum condition and analyticity in coordinate space in the axiomatic domain.

It is nevertheless instructive to have a look. Suppose that some x_{ij} in expression (50) is positive time-like. Then

$$(-\frac{1}{2}x_{ij}^2 + i\epsilon x_{ij}^0)^{-\delta_{ij}} = (\frac{1}{2}|x_{ij}^2|)^{-\delta_{ij}} e^{(\pi-\epsilon)\Im m\ \delta_{ij}} \cdot phase$$

The second factor increases exponentially when the imaginary part of δ_{ij} becomes large. The Γ-functions in expression (50) decay exponentially in the imaginary directions. One finds in a case by case study for $n = 4$ that the Γ-functions always keep the integrand bounded at large imaginary parts of δ_{ij}, provided M^c does not grow exponentially.

3.5 Time ordered Green functions in Minkowski space

The time ordered Green functions are vacuum expectation values of time ordered products of fields. Time ordered means that the fields with later

times as arguments stand further to the left. According to this definition, they can be expressed in terms of the Wightman functions. For Bose fields

$$\langle \Omega, T\left\{\phi^{i_n}(x_n)...\phi^{i_1}(x_1)\right\}\Omega\rangle = \langle \Omega, \phi^{i_{\pi n}}(x_{\pi n})...\phi^{i_{\pi 1}}(x_{\pi 1})\Omega\rangle \tag{55}$$

where π is that permutation of $1, ..., n$ which assures that

$$x^0_{\pi n} \geq ... \geq x^0_{\pi 2} \geq x^0_{\pi 1} \ .$$

Inserting the Mellin representation of the Wightman function and using the locality property (49) of the Mellin amplitude, it follows that the time ordered Green function in Minkowski space possesses the Mellin representation

$$\begin{aligned} &\langle \Omega, T\left\{\phi^{i_n}(x_n)...\phi^{i_1}(x_1)\right\}\Omega\rangle \\ &= (2\pi i)^{-n(n-3)/2}\int\int d\delta\ M^c_{i_n...i_1}(\{\delta_{ij}\}) \\ &\qquad \prod_{i>j}\Gamma(\delta_{ij})(-\frac{1}{2}x^2_{ij}+i\epsilon)^{-\delta_{ij}}. \end{aligned} \tag{56}$$

It differs from the Wightman functions only through the $i\epsilon$-prescriptions.

$\Gamma(\delta)(-\frac{1}{2}x^2+i\epsilon)^{-\delta}$ is a generalized Feynman propagator in the sense of Speer [66]. Therefore the x-dependent factor is the inverse Fourier-transform of a generalized Feynman integral. In momentum space, the result looks like expression (53), (54) except that the factor $\rho_D(\delta)$ is different, p_{ij}-integrations are over all of momentum space, and p^2 is to be read as $p^2+i\epsilon$, as is appropriate for Feynman propagators.

4 Relations to theories in lower dimensions

4.1 Dimensional reduction: AdS-orbits & their boundary

The conformal group in D dimensions, G = universal covering of SO(2,D) , acts transitively on the tube $\mathcal{M}_D \simeq S^{D-1}\times \mathbf{R}$ (the ∞-sheeted cover of conformally compactified Minkowski space). Under the subgroup which covers $SO(2, D-1)$, $\mathcal{M}_D$ decomposes into orbits as follows

$$\mathcal{M}_D = \mathcal{M}_{D-1} \cup AdS_D \cup AdS_D \tag{57}$$

There are two copies of Anti-de Sitter space in D dimensions - or rather, its simply connected covering. The first factor is their common boundary. Points on the space $\mathcal{M}_D$ can be coordinatized by (e, τ), $\infty < \tau < \infty$, and e a unit D-vector with components $(e_1, ..., e_{D-1}, e_{D+1})$ The orbit $\mathcal{M}_{D-1}$ consists of points with $e_{D-1} = 0$, the equator of the sphere S^{D-1}.

This is most easily seen by using Dirac's manifestly covariant formalism [10]. In place of points x of (conformally compactified) Minkowski space it uses rays of light-like $D+2$-vectors $\xi = (\xi_0, \xi_1 ... \xi_{D-1}, \xi_{D+1}, \xi_{D+2})$. Light-like means

$$(\xi_0)^2 - (\xi_1)^2 - ... - (\xi_{D-1})^2 - (\xi_{D+1})^2 + (\xi_{D+2})^2 = 0 \tag{58}$$

Vectors ξ and $\xi' = \lambda\xi$, $\lambda > 0$ are on the same ray.

Elements of the group $SO(2, D)$ act as pseudo-rotations on the $D+2$-vectors ξ. The subgroup $SO(2, D-1)$ leaves ξ_{D-1} invariant. There are therefore two invariant domains: rays with $\xi_{D-1} = 0$ and rays with $\xi_{D-1} \neq 0$. The covering of the first domain is $\mathcal{M}_{D-1}$. For the second domain we may use homogeneity to scale ξ to $r\xi$, $r > 0$ such that either $\xi_{D-1} = 1$ or $\xi_{D-1} = -1$. In either case Eq.(58) then tells us that we are dealing with Anti-de Sitter space.

The relation between the rays and the coordinates (τ, e) is as follows [49]:

$$\xi_0 = r\cos\tau,\ \xi_{D+2} = r\sin\tau,\ \xi_k = re_k \tag{59}$$

for $(k = 1, ..., D-1, D+1)$. Therefore, $\xi_{D-1} = 0$ means $e_{D-1} = 0$ as claimed, and $\xi_{D-1} > 0$ means $e_{D-1} > 0$ while $\xi_{D-1} < 0$ means $e_{D-1} < 0$.

For future use I also give the relation between Minkowski space coordinates x_μ and ξ. Set

$\kappa = \xi_{D+1} + \xi_{D+2}$. Then $x_\mu = \xi_\mu/\kappa, \quad (\mu = 0...D-1)$.

Abbreviating $\{i_n, ..., i_1\} = a$ as before, the Mellin representation of a conformally invariant scalar n-point Wightman function can be written in manifestly covariant form as follows.

$$\begin{aligned} &W_a(\xi^n, ..., \xi^1) \equiv \kappa_1^{-d_1} ... \kappa_n^{-d_n} W_a(x_n, ..., x_1) \\ &= (2\pi i)^{-n(n-3)/2} \int d\delta M_a(\{\delta_{ij}\}) \prod_{i>j} (\xi_i \xi_j)^{-\delta_{ij}} \ . \end{aligned} \tag{60}$$

We see that the restriction of the amplitude W_a to $\mathcal{M}_{D-1} \times ... \times \mathcal{M}_{D-1}$, (i. e to points with $\xi^i_{D-1} = 0$) exists and is given by identically the same Mellin representation with the same Mellin amplitude M_a, with the understanding

that $\xi^i_{D-1} = 0$. Conversely, this restriction suffices to determine the Mellin amplitude M_a, and the Mellin amplitude yields the amplitude W_a on all of $\mathcal{M}_D \times ... \times \mathcal{M}_D$. This is why I call the Mellin representation a *D-independent representation*.

Wightman positivity as discussed in section 2.2 can be formulated for scalar amplitudes $W_a(\xi_n, ...\xi_1)$ by using sequences of test functions

$$f^{\emptyset}, f^{i_1}(\xi^1)...f^{i_1...i_N}(\xi^1, ..., \xi^N) \tag{61}$$

which are homogeneous functions of ξ_k of degree $-D + d_k$. The existence of the restriction implies that test functions are permissible which are concentrated on the hyper-surface $\xi^k_{D-1} = 0$, i.e.

$$f^{i_1...i_k}(\xi^1, ..., \xi^k) \propto \delta(\xi^1_{D-1})...\delta(\xi^k_{D-1}) \tag{62}$$

Thus, Wightman positivity of the restricted Wightman functions in $D-1$ dimensions follows from Wightman positivity of the original Wightman functions. Therefore *a unitary CFT in D dimensions restricts to a unitary CFT in* $D-1$ *dimensions*.

The operator content of $D-1$-dimensional CFT's which are obtainable by dimensional reduction has special properties. Together with every field ϕ^i of dimension d_i there is a tower of fields $\phi^i_{,n}$, $n = 1, 2, 3, ...$ of dimensions $d_i + n$. They arise because derivatives of fields in D dimensions need not be derivatives in $D-1$ dimensions. In the manifestly covariant formalism,

$$\phi^i_{,n} = D_{D-1}...D_{D-1}\phi^i|_{\xi_{D-1}=0} \tag{63}$$

(n factors D_{D-1}). D_A is the $SO(2, D)$ covariant interior differential operator on the $D+2$-dimensional cone $\xi^2 = 0$ [5]. The missing generators J_{AB}, $B = D-1$ of $SO(2, D)$ transform $\phi^i \leftrightarrow D_{D-1}\phi^i$, leaving 2- and 3-point functions invariant.

4.2 Dual amplitudes from 2-dimensional conformal field theories

In two dimensions, some things are different. $S^1 \times \mathbf{R}$ is not simply connected, and is therefore not the universal covering of compactified Minkowski space. Spin needs not be half integral, and more general types of statistics, known as braid statistics, are possible [24]. There exist chiral theories, in which

fields depend only on one of the light-cone variables $x^{\pm} = x^0 \pm x^1$; they are effectively 1-dimensional. All this gives rise to possibilities which do not exist in higher dimensions, such as anyons [2].

Let us examine sufficient conditions, such that a universal Mellin representation as discussed in section 3 exists and furnishes a dual amplitude. It will turn out that invariance under space reflections and conventional locality, i.e. Bose/Fermi-statistics are sufficient.

In two dimensionsional Minkowski space $x^2 = x^+x^-$. As a result, the anharmonic ratios factorize,

$$\omega_1 = \omega_1^+\omega_1^-, \quad \omega_2 = \omega_2^+\omega_2^-, \quad \omega_3 = \omega_3^+\omega_3^-, \tag{64}$$

$$\omega_1^{\pm} = \frac{x_{12}^{\pm}x_{34}^{\pm}}{x_{13}^{\pm}x_{24}^{\pm}}, \quad \omega_2^{\pm} = \frac{x_{14}^{\pm}x_{23}^{\pm}}{x_{12}^{\pm}x_{34}^{\pm}}, \quad \omega_3^{\pm} = \frac{x_{13}^{\pm}x_{24}^{\pm}}{x_{14}^{\pm}x_{23}^{\pm}}. \tag{65}$$

They share the property that $\omega_1\omega_2\omega_3 = 1$. But it follows from the definition (65) that the three quantities ω_i^+ are actually all dependent. Take ω_1^+ as the independent one and call it x, temporarily. Then

$$\omega_1^+ = x, \quad (\omega_3^+)^{-1} = 1 - x, \quad \omega_2^+ = (1-x)/x, \tag{66}$$

and similarly for ω_i^-.

Space reflection takes $x^1 \mapsto -x^1$, therefore $x^+ \leftrightarrow x^-$ and $\omega_i^+ \leftrightarrow \omega_i^-$. Let us now use the above equations to determine the two independent quantities ω_1^+ and ω_1^- in terms of two independent anharmonic rations ω_1 and ω_2 of the two-dimensional theory. There are two solutions which go into each other by interchanging $\omega_1^+ \leftrightarrow \omega_1^-$. This is expected since space reflections leave all ω_i invariant, but interchange ω_1^+ and ω_1^-. In other words, ω^+ and ω^- depend on a variable $\sigma = \pm 1$ in addition to ω_1, ω_2. Explicitly,

$$\begin{aligned} \omega^+ &= -b + \sigma\sqrt{b^2 - \omega_1}, \\ \omega^- &= -b - \sigma\sqrt{b^2 - \omega_1}, \\ b &= -\frac{1}{2}\left(\omega_1(\omega_2 - 1) - 1\right) \ . \end{aligned} \tag{67}$$

Retain the notation ω for triples $(\omega_1, \omega_2, \omega_3)$ subject to $\omega_1\omega_2\omega_3 = 1$. Conformal 4-point Green functions $G_{i_4\ldots i_1}$ are products of kinematical factors times functions of ω_1^+ and ω_1^-, that is functions $F_{i_4\ldots i_1}(\omega, \sigma)$.

Under space reflections, $\sigma \mapsto -\sigma$, while ω is invariant. Suppose now that the 2-dimensional theory is space reflection invariant and obeys conventional locality. Then $F_{i_4\ldots i_1}(\omega, \sigma)$ is independent of σ, and obeys the crossing relation (37).

Now we may Mellin transform $F_{i_4 \ldots i_1}(\omega)$ as in eq.(42) and this yields a dual amplitude $\hat{M}^c(\beta)$.

In terms of the quantities $\omega_1^{\pm}$, the factor

$$\begin{aligned} \omega_1^{-s_1}\omega_2^{-s_2}\omega_3^{-s_3} = \quad & (\omega_1^+)^{-\beta_{12}}(1-\omega_1^+)^{-\beta_{23}} \\ & (\omega_1^-)^{-\beta_{12}}(1-\omega_1^-)^{-\beta_{23}} \ . \end{aligned} \tag{68}$$

Space reflection invariance is not a necessary condition. Consider chiral theories whose fields depend only on one variable; let us say x^+. The 4-point amplitudes will be equal to kinematical factors times functions $F_{i_4 \ldots i_1}(\omega^+)$ of anharmonic ratios. There is now only one independent anharmonic ratio, and Mellin transformation is of no help. But suppose that the construction of F yields an integral representation

$$\begin{aligned} F_a(\omega^+) \ &= \ (2\pi i)^{-2} \int\int d\beta_{12} d\beta_{23} \hat{M}_a(\{\beta_{ij}\}) \\ & \qquad (\omega_1^+)^{-\beta_{12}}(1-\omega_1^+)^{-\beta_{23}}. \end{aligned} \tag{69}$$

Then this can serve as a substitute for a Mellin transformation. Considering $\hat{M}_a$ as a function of variables $\beta_{ij} = \beta_{ji}$, $i \neq j$ subject to the constraints (40,41) will serve to define the action $\beta \mapsto \pi\beta$ of permutations π on $\beta = \{\beta_{ij}\}$. Suppose that $F_a(\omega)$, $a = i_4, ..., i_1$ is invariant under the fractional linear transformations of ω_1^+ which amount to permutations π of ω_i^+ as defined by eq.(66), invariance meaning that $F_{\pi a}(\pi\omega^+) = F_a(\omega^+)$. Suppose that this is implemented by $\hat{M}_{\pi a}(\pi\beta) = \hat{M}_a(\beta)$. And suppose further that $\hat{M}_a(\{\beta_{ij}\})$ has poles in individual variables, with residues that are polynomials in the other(s). Then $\hat{M}_a$ furnishes a dual amplitude.

When F_a come from a dual resonance model, then the supposition regarding poles and residues is true by construction.

5 Euklidean partial wave expansion of scalar 4-point functions

5.1 Fundamental fields

Fields with dimension $d < \frac{D}{2}$ will be called *fundamental fields*. By the restrictions on χ, they must be either scalar or spinor fields, in $D = 4$ dimensions. A theory may have fundamental fields or not. This does not affect

duality. But fundamental fields play a special role in the Euklidean partial wave expansions, requiring exorcision of a ghost - the shadow pole of Ferrara Gatto, Grillo and Parisi [16]. Absence of this ghost is equivalent to validity of renormalized Schwinger Dyson equations for all n-point Green functions in Lagrangean field theory, see the end of subsection 5.2.

5.2 Euklidean partial wave expansions

Typically, the scalar Euklidean Greens functions are correlation functions of a statistical mechanical system, e.g.

$$G_{i_4 i_3 i_2 i_1}(\mathbf{x}_4, ..., \mathbf{x}_1) = \\ < \varphi^{Ei_4}(\mathbf{x}_4), \varphi^{Ei_3}(\mathbf{x}_3), \varphi^{Ei_2}(\mathbf{x}_2), \varphi^{Ei_1}(\mathbf{x}_1) > .$$

where the (commuting) Euklidean fields φ^E are variables of the statistical mechanical system. Suppose that we replace the two fields on the right, $\varphi^{E2}(\mathbf{x}_2)$ and $\varphi^{E1}(\mathbf{x}_1)$ by their conformally transformed brothers, transformed by one and the same element $g \in G^E$. The result will be a function $\hat{G}_{i_4 i_3 i_2 i_1}$ of g. It follows from the transitivity of the action of the Euklidean conformal group on triples of distinct points, that G_{0000} is determined for all $\mathbf{x}_1, ..., \mathbf{x}_4$ if we know this function $\hat{G}$ of g for some standard choice of $\hat{\mathbf{x}}_1, ..., \hat{\mathbf{x}}_4$, with $\hat{\mathbf{x}}_1 \neq \hat{\mathbf{x}}_2, \hat{\mathbf{x}}_3 \neq \hat{\mathbf{x}}_4$. One may now perform a partial wave expansion (i.e. expansion in representation functions of G^E of $\hat{G}(g)$ on G^E) for this standard choice of $\hat{\mathbf{x}}_4, ..., \hat{\mathbf{x}}_1$. For scalar amplitudes, such partial wave expansions have been extensively studied in the 70's. The program was initiated by the author [42, 43, 44] and completed by Todorov and his collaborators [14]. Normalization factors are computed in [11]; a review in book form is [12]. The result, rewritten for $G_{i_4 i_3 i_2 i_1}$ is as follows.

It is convenient to introduce the notation

$$h = \frac{D}{2} \tag{70}$$

The representations which enter are labeled [4] by $\chi = [l, \delta = h + c]$ just like the representations of G, where l is now interpreted as a representation of the Euklidean Lorentz group $M^E = Spin(D)$ in a vector space V^l (this makes no difference by Weyl's unitary trick). But the dimension δ is now either of

[4] The representations which we here label by $\chi = [l, h + c]$ are labeled by $\chi = [l, c]$ in the literature on Euklidean partial wave expansions

the form $\delta = h + c$, c imaginary (*principal series of representations*) or c real in some interval (*complementary series*). There is also a *discrete series* if D is odd, but it does not contribute to scalar amplitudes. The representations $\chi = [l, h + c]$ and $\tilde{\chi} = [\bar{l}, h - c]$ of the principal or complementary series are equivalent, where $\bar{l}$ is the dual representation to l; for symmetric tensor representations $\bar{l} \simeq l$. I write $l = 0$ for the trivial representation.

The unitary representations of G^E are special cases of elementary representations. They are constructed as induced representations and are labeled by $\chi = [l, h + c]$ with complex c.

Suppose the scalar Euklidean fields $\varphi^{Ei_1}, ...$ have real dimensions $d_1, ..., d_4$ so that they transform according to representations $\chi_1 = [0, d_1], ...\chi_4 = [0, d_4]$. Given $\chi = [l, \delta]$, a conformal invariant three point function $\mathfrak{V}(\mathbf{x}_3, \chi, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) \in V^l$ exists only if l is a traceless symmetric tensor representation of M^E; in this case it is unique up to normalization.

There exists a unique canonically normalized conformal invariant 2-point function $\Delta^\chi(\mathbf{x}, \mathbf{y})$. The three point functions can be normalized in such a way that

$$\begin{aligned} &\int d^D\mathbf{y} \Delta^\chi(\mathbf{x}, \mathbf{y}) \mathfrak{V}(\mathbf{y}, \tilde{\chi}, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) \\ &= \mathfrak{V}(\mathbf{x}, \chi, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) \end{aligned} \tag{71}$$

Validity of this amputation identity for arbitrary χ requires that

$$\Delta^{\tilde{\chi}} = (\Delta^\chi)^{-1} \ , \tag{72}$$

the inverse being meant in the convolution sense.

Referring to 4-point functions only for now, a *channel* is a division of the four arguments of a 4-point function labeled by $4, 3, 2, 1$ into two groups of two, viz. (43)(21) or (42)(31) or (41)(32). I will sometimes write (21) in place of (43)(21) for short.

There is no way of doing partial wave expansions of 4-point Wightman functions in positive energy representations in different channels without commuting the fields. But for Euklidean Green functions this can be done, because they are symmetric in their arguments - the classical statistical mechanical observables $\varphi^{Ei_k}(\mathbf{x}_k)$ commute - in the bosonic case. The above mentioned expansion is the partial wave expansion in the channel (43)(21).

The Euklidean partial wave expansion of the *connected* scalar 4-point function of four hermitean scalar field $\phi^{i_4}, ..., \phi^{i_1}$ with spin and dimension

$\chi_k = [0, d_k], k = 4, ..., 1$, in a theory in which no fundamental field appears in the OPE of both $\phi^{i_2}\phi^{i_1}$ and $\phi^{i_3}\phi^{i_4}$ reads as follows

$$G^c_{i_4 i_3 i_2 i_1}(\mathbf{x}_4, \mathbf{x}_3, \mathbf{x}_2, \mathbf{x}_1) = \int d\chi g^{(43)(21)}_{i_4 i_3 i_2 i_1}(\chi) \tag{73}$$
$$\int d^D\mathbf{x} \langle \mathfrak{V}(\mathbf{x}, \tilde{\chi}, \mathbf{x}_4, \chi_4, \mathbf{x}_3, \chi_3), \mathfrak{V}(\mathbf{x}, \chi, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1)\rangle$$

The superscript (43)(21) on $g(\chi)$ indicates the channel in which we expand. $\langle , \rangle$ is contraction of tensor indices.

The χ-integration is over the part of the principal series $\chi = [l, \frac{D}{2} + c]$, c imaginary, where l is a completely symmetric traceless tensor representation, with Plancherel measure $d\chi$. Thus [14]

$$\begin{aligned} \int d\chi ... &= \frac{1}{2\pi i} \sum_l \int_{-i\infty}^{\infty} dc\ \rho_l(c) ... \\ \rho_l(c) &= \frac{\Gamma(l+h)}{2(2\pi)^h l!} \frac{\Gamma(h-1+c)\Gamma(h-1-c)}{\Gamma(c)\Gamma(-c)} \\ &\quad [(h+l-1)^2 - c^2] \end{aligned} \tag{74}$$

In even dimensions D, the Plancherel weight $\rho_l(c)$ is a polynomial in c. Expansion (73) can be written in a more symmetrical form by inserting eq.(71), but is more convenient for our purpose as it stands.

The contribution from a single χ is called the partial wave. Since all the other factors are kinematically determined, I will also refer to $g^{(43)(21)}$ as the partial wave. It depends on normalization conventions including (71).

If the OPE of both $\phi^{i_2}\phi^{i_1}$ and $\phi^{i_3}\phi^{i_4}$ contains a fundamental scalar field of dimension d_f then a Born term must be added to the right hand side of eq.(73). It has the same form as the integrand of the χ-integration, with $\chi_f = [0, d_f]$ substituted for χ, and a product of coupling constants substituting for $g^{(43)(21}_{i_4 i_3 i_2 i_1}(\chi)$. It can be included by altering the path of the c-integration, see section 6 below and figure 1 a).

Let us now consider situations in which there exist non-vanishing disconnected parts other than a contribution $G_{i_4 i_3}(\mathbf{x}_4, \mathbf{x}_3) G_{i_2 i_1}(\mathbf{x}_2, \mathbf{x}_1)$ [which may be interpreted as the contribution of the unit operator in the OPE]. This will be so in the case of four equal hermitean scalar fields $\phi^{i_1} = \phi^{i_2} = \phi^{i_3} = \phi^{i_4} = \phi^0$ with dimension d_0. Set $\chi_0 = [0, d_0]$.

There exists a choice of normalization factors such that the coefficient in the Euklidean partial wave expansion of the sum $G_{00}(\mathbf{x}_4, \mathbf{x}_2) G_{00}(\mathbf{x}_3, \mathbf{x}_1) +$



Figure 1: Path of the c-integration in the complex c-plane in the derivation of OPE, for fixed l in the presence of a fundamental field with Lorentz spin l and dimension $d_f = h + c_f$, $c_f < 0$: a) before and b) after the shift of the path of integration. To each field ϕ^k in the OPE of Lorentz spin l and dimension $d_k = h + c_k$ there corresponds a pole $\bullet$ at $c = c_k$ in $g^{(43)(21)}_{\dots}([l, c])$, and a shadow pole $\circ$ at $c = -c_k$. The contributions in a) from the the circles around the poles at $c = \pm c_f$ represent the Born term.

$G_{00}(\mathbf{x}_4, \mathbf{x}_1)G_{00}(\mathbf{x}_3, \mathbf{x}_2)$ of disconnected parts is identically equal to 1. This will be the case if the normalization factors in eq.(76) for $\mathfrak{V}$ satisfy eq.(79).

Adopting these normalization conventions, the expansion of the full Green function will be

$$\begin{aligned} & G^4_{0000}(\mathbf{x}_4, \mathbf{x}_3, \mathbf{x}_2, \mathbf{x}_1) \qquad (75) \\ = \;& G_{00}(\mathbf{x}_4, \mathbf{x}_3)G_{00}(\mathbf{x}_2, \mathbf{x}_1) + \\ + \;& \int d\chi[1 + g^{(43)(21)}_{0000}(\chi)] \int d^D\mathbf{x} \\ & \quad \langle \mathfrak{V}(\mathbf{x}, \tilde{\chi}, \mathbf{x}_4, \chi_0, \mathbf{x}_3, \chi_0), \mathfrak{V}(\mathbf{x}, \chi, \mathbf{x}_2, \chi_0, \mathbf{x}_1, \chi_0)\rangle \\ + \;& \text{ possibly a Born term} \end{aligned}$$

The representation (75) is also valid for a generalized free field theory, with $g^{(43)(21)}(\chi) = 0$.

5.3 Explicit form of the 3-point function

In our later computations, we will need the explicit form of the three point function $\mathfrak{V}(\mathbf{x}, \chi, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1)$.

It is convenient for our purposes to use a particular realization of the representation space V^l of traceless symmetric tensors of rank l. With every such tensor $t_{\mu_1,\ldots,\mu_l}$ one associates a homogeneous polynomial in a complex D-vector $\mathfrak{z}$ with $\mathfrak{z}^2 = 0$,

$$t(\mathfrak{z}) = \sum t_{\mu_1,\ldots,\mu_l} \mathfrak{z}^{\mu_1} \ldots \mathfrak{z}^{\mu_l},$$

and one regards V^l as a space of such homogeneous polynomials. [Here and everywhere, I simply write l for the rank of a symmetric tensor representation l].

In this realization, the vertex function $\mathfrak{V}$ become functions of $\mathfrak{z}$.

I use notations for the normalization factors which are in accord with the literature [14]. Let $d_i = h + c_i$ and set $c^+ = \frac{1}{2}(c_1 + c_2)$, $c_- = \frac{1}{2}(c_1 - c_2)$ and $\chi = [l, \delta] = [l, h + c]$. Then

$$\begin{aligned} \mathfrak{V}(\mathbf{x}_0, \chi, \mathfrak{z}, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) &= N_l(c_+, c_-, c) \\ &\quad \mathfrak{V}^u(\mathbf{x}_0, \chi, \mathfrak{z}, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) \end{aligned} \tag{76}$$

with

$$\begin{aligned} &\mathfrak{V}^u(\mathbf{x}_0, \chi, \mathfrak{z}, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) = (2\pi)^{-h} (\lambda \cdot \mathfrak{z})^l \\ &\left(\frac{2}{\mathbf{x}_{12}^2}\right)^{\frac{1}{2}(h-c+l)+c_+} \left(\frac{4}{\mathbf{x}_{10}^2 \mathbf{x}_{20}^2}\right)^{\frac{1}{2}(h+c-l)} \left(\frac{\mathbf{x}_{20}^2}{\mathbf{x}_{10}^2}\right)^{c_-} , \end{aligned} \tag{77}$$

$$\lambda = 2\left(\frac{\mathbf{x}_{10}}{\mathbf{x}_{10}^2} - \frac{\mathbf{x}_{20}}{\mathbf{x}_{20}^2}\right). \tag{78}$$

The right hand side defines the analytic continuation throughout the axiomatic analyticity domain. In Euklidean space $-x^2 = \mathbf{x}^2 > 0$. In Minkowski space, $i\epsilon$-prescriptions have to be put , which depend on the ordering of the fields, similarly as in section 3.4.

There exists a canonical choice of the normalization factors $N_l(c_+, 0, c)$ in the expression (76) for the 3-point function . They have been determined in the literature from the requirement that appropriate orthonormality properties of the expansion functions, and amputation identities such as eq.(71)

hold [11]. The result gives the product

$$N_l(c_+,0,c)N_l(c_+,0,-c) = \qquad (79)$$
$$\frac{\Gamma(\frac{1}{2}(h-c+l)+c_+)\Gamma(\frac{1}{2}(h+c+l)+c_+)}{\Gamma(\frac{1}{2}(h-c+l)-c_+)\Gamma(\frac{1}{2}(h+c+l)-c_+)}$$

One sees that this is a meromorphic function of c.

5.4 Note on the use of normalization conventions

Let us finally comment on normalization factors in the general case of not necessarily equal dimensions d_4, d_3, d_2, d_1 of the external scalar fields.

Disconnected parts other than the first term in the Euklidean partial wave expansion (75) can only appear when either $d_1 = d_3$ and $d_2 = d_4$ or $d_1 = d_4$ and $d_2 = d_3$. When neither is the case, it is less convenient to use the canonical normalization factors, because they have branch cuts such that the product $N_l(c_+^{12}, c_-^{12}, c)N_l(c_+^{34}, c_-^{34}, -c)$ is in general not meromorphic. One may choose $N_l \equiv 1$ instead, at the cost of introducing (meromorphic) factors in amputation identities like eq.(71). The orthogonality relations remain unchanged. In any case, the quantity

$$N_l(c_+^{12}, c_-^{12}, c)N_l(c_+^{34}, c_-^{34}, -c)g_{i_4i_3i_2i_1}^{(43)(21)}([l, h+c]) \qquad (80)$$

is independent of normalization conventions, and validity of OPE requires meromorphy of this product.

In the case of equal dimensions $d_4, ..., d_1$ the canonical product (79) of normalization factors is meromorphic, so that also $g^{(43)(21)}([l, h+c])$ is meromorphic.

5.5 OPE from Euklidean partial wave expansions

From eq.(75), operator product expansions (19) for the 4-point function have been derived, assuming meromorphy properties of partial waves $g^{(43)(21)}([l, c])$. A complete derivation was given in ref. [14] for even dimension D and under some restrictions on the field dimensions $d_1, ..., d_4$. It was stated that the result can be generalized. I will briefly review the result and slightly generalize it to emphasize factorization.

The first step exploits the equivalence of representations $\chi = [l, h+c]$ and $\tilde{\chi} = [l, h-c]$ of the Euklidean conformal group to split the three point

function into Clebsch-Gordan kernels of the second kind in a way modeled after the split of Legendre functions P_l into Legendre functions of the second kind, Q_l,

$$\mathfrak{V}(\mathbf{x}_0, \tilde{\chi}, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) = \tag{81}$$
$$\frac{\pi}{\sin(l+c)}[\mathfrak{Q}(\mathbf{x}, \tilde{\chi}, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) -$$
$$- \int d^D\mathbf{y} \Delta^{\tilde{\chi}}(\mathbf{x}, \mathbf{y}) \mathfrak{Q}(\mathbf{y}, \chi, \mathbf{x}_2, \chi_1, \mathbf{x}_1, \chi_2)] \ .$$

such that the partial Fourier transform $\mathfrak{Q}(p, \tilde{\chi}, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1)$ is an entire holomorphic function of p.

As a consequence of this split, the integrand of the Euklidean partial wave expansion becomes a sum of two terms. Upon a change of variables $c \mapsto -c$, the second term becomes equal to the first, producing a factor of 2.

The result can be rewritten in terms of the partial Euklidean Fourier transforms as

$$G^c_{i_4 i_3 i_2 i_1}(\mathbf{x}_4, \mathbf{x}_3, \mathbf{x}_2, \mathbf{x}_1) = \int d\chi \frac{2\pi}{\sin \pi(l+c)} g^{(43)(21)}_{i_4 i_3 i_2 i_1}(\chi)$$
$$\int (d\mathbf{p}) \langle \mathfrak{Q}(-p, \tilde{\chi}, \mathbf{x}_4, \chi_4, \mathbf{x}_3, \chi_3), \mathfrak{V}(p, \chi, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1) \rangle$$

Integration is over Euklidean momenta, i.e. over imaginary $p^0 = ip^D$, with $(d\mathbf{p}) = (2\pi)^{-D} dp^1 ... dp^D$.

Let us now restrict attention to (Euklidean) arguments $\mathbf{x}_i$ where the Euklidean time components satisfy inequalities

$$x_1^D > 0, \ x_2^D > 0 \ , \ x_3^D < 0, \ x_4^D < 0 \ . \tag{82}$$

Inserting the split (81) with χ substituted for $\tilde{\chi}$ also for the remaining factor $\mathfrak{V}$, a sum of two terms appears, but the term involving the entire function $\mathfrak{Q}(-p, \tilde{\chi}...)\mathfrak{Q}(p, \chi, ...)$ of p will make zero contribution after the following deformation of the path of the p-integration (figure 2. The path of the p^0-integration is deformed as shown in figure 2. The closure of the path is allowed, because the integrand has exponential falloff as $\Re e \ p^0 \mapsto \infty$.

Let $\Delta^\chi(p)$ be the function of p which is the analytic continuation of the Euklidean Fourier transform of Δ^χ, with a cut along the positive p^0-axis. Writing the discontinuity as

$$\frac{i}{\pi} \sin \pi(l-c) \Delta^\chi_+(p) \ ,$$

then $\Delta_+^\chi(p)$ is a Wightman function which is positive when χ labels a positive energy representation of the conformal group.



Figure 2: Deformation of the path of integration in the complex energy plane. a) Integration over Euklidean space involves integration over imaginary p^0. b) The deformed path

In the light of the remarks of the last subsection, I will extract normalization factors,

$$\begin{aligned}\mathfrak{Q}(\mathbf{x},\tilde{\chi},\mathbf{x}_2,\chi_2,\mathbf{x}_1,\chi_1) &= N_l(c_+^{12},c_-^{12},-c)\\ &\quad \mathfrak{Q}^u(\mathbf{x},\tilde{\chi},\mathbf{x}_2,\chi_2,\mathbf{x}_1,\chi_1)\end{aligned}$$

An explicit formula for $\mathfrak{Q}^u$ is given the appendix, eq.(175).

The notation for the normalization factor has been chosen to be in agreement with the literature [14]; $c_\pm^{21} = \frac{1}{2}(c_2 \pm c_1)$ when $d_1 = h + c_1$, $d_2 = h + c_2$.

Extracting normalization factors, the result after the deformation of the

path of integration reads

$$G^c_{i_4i_3i_2i_1}(\mathbf{x}_4,\mathbf{x}_3,\mathbf{x}_2,\mathbf{x}_1) = \int d\chi \frac{-2\pi}{\sin\pi(l+c)}$$
$$N_l(c^{34}_+,c^{34}_-,-c)N_l(c^{12}_+,c^{12}_-,c)g^{(43)(21)}_{i_4i_3i_2i_1}(\chi)$$
$$\int_{V_+}(dp)\langle\mathfrak{Q}^u(-p,\tilde{\chi},\mathbf{x}_4,\chi_4,\mathbf{x}_3,\chi_3),$$
$$\Delta^\chi_+(p)\mathfrak{Q}^u(p,\tilde{\chi},\mathbf{x}_2,\chi_2,\mathbf{x}_1,\chi_1)\rangle$$

The integration now runs over the forward light cone V_+ in Minkowskian momentum space, $(dp) = (2\pi)^{-D}dp^0...dp^{D-1}$.

Thereafter, the path of the c-integration is closed in the half plane $Re\ c > 0$ as shown in figure 1, assuming growth conditions on partial wave amplitudes $g(\chi)$. Remember that $d\chi$ includes a factor $\rho_l(c)$, the Plancherel weight. Assume meromorphy of

$$\rho_l(c)\ N_l(c^{34}_+,c^{34}_-,-c)N_l(c^{12}_+,c^{12}_-,c)\ g^{(43)(21)}_{i_4i_3i_2i_1}(\chi) \tag{83}$$

in c with simple poles at real positions $c = c_k > 0$ for $l = l_k$. Every such pole will make a discrete contribution which is in accord with the presence of a field ϕ^k in the OPE with spin and dimension $l_k, d_k = h + c_k$, *provided* the residue has factorization properties as detailed below. The absence of contributions from other poles, notably from the poles of $\pi/\sin\pi(l+c)$ in even dimensions D is a subtle issue in the derivation of OPE. It was discussed in detail in [14] and will concern us again later. In the present paper, I assume the validity of OPE and conclude that the partial waves $g(\chi)$ must have properties such that there are no other contributions, cf. section 6.2.

Suppose expression (83) has simple poles at $\chi = \chi_k = [l_k, h + c_k]$ with residues which factorize as follows

$$\frac{2\pi}{\sin\pi(c_k+l)} res_{c=c_k}$$
$$\left[\rho_l(c)N_l(c^{34}_+,c^{34}_-,-c)N_l(c^{12}_+,c^{12}_-,c)\ g^{(43)(21)}([l,h+c])\right]$$
$$= \bar{g}^{i_3i_4}_k g^{i_2i_1}_k\ . \tag{84}$$

Then we obtain as our final result

$$G_{i_4i_3i_2i_1}(\mathbf{x}_4...\mathbf{x}_1) = \sum_k \bar{g}^{i_3i_4}_k g^{i_2i_1}_k \int_{V_+}(dp) \tag{85}$$
$$\langle\mathfrak{Q}^u(-p,\tilde{\chi}_k,\mathbf{x}_2,\chi_2,\mathbf{x}_1,\chi_1)\Delta^\chi_+(p)\mathfrak{Q}^u(p,\tilde{\chi}_k,\mathbf{x}_4,\chi_4,\mathbf{x}_3,\chi_3)\rangle$$

This admits analytic continuation to Minkowski space, and satisfies Osterwalder Schrader positivity as shown below.

Consider now also the situation with disconnected parts. Eq.(85) is valid as it stands for G_{0000} except that the contribution $G_{00}(\mathbf{x}_4\mathbf{x}_3)G_{00}(\mathbf{x}_2\mathbf{x}_1)$ of the unit operator to the OPE must be added on the right hand side. But in the definition (84) of coupling constants, $[1 + g_{0000}^{(43)(21)}(\chi)]$ is to be substituted for $g_{i_4i_3i_2i_1}^{(43)(21)}(\chi)$, and normalization conventions (79) are in force.

In conclusion, there is a bijective correspondence between poles at $\chi = \chi_k = [l_k, h + c_k],\ c_k > 0$ of expression (83) and contributions of fields ϕ^k with conformal transformation law specified by χ_k to the OPE for the 4-point function, and the residues must have appropriate factorization properties.

If there is a fundamental field ϕ^f transforming according to $\chi_f = [0, d_f]$ with $d_f < D/2$ in the OPE then the above statement must be qualified as follows: Among the poles of expression (83), there is a pole at $\chi = \chi_f$ whose contribution must be included, but the contribution of its "shadow" at $\chi = \tilde{\chi_f} = [l, D - d_f]$ must be omitted, as shown in figure 1 b).

This has a deep reason which was extensively discussed in the literature [42, 43, 14], and will be briefly reviewed in section 6 below.

5.6 Osterwalder Schrader positivity

In section 2 it was pointed out that validity of OPE implies positivity of Wightman functions. This statement has a Euklidean version. It involves Osterwalder Schrader (OS) positivity, also known as reflection positivity [29]. This result is useful for us because it saves us from having to worry about validity of the spectrum condition. Validity of the spectrum condition follows from OS- positivity and other Euklidean axioms by the Osterwalder Schrader reconstruction theorem [57, 58]. The other Euklidean axioms include crossing symmetry and are fulfilled by construction in the present approach.

Similarly as for Wightman functions, it suffices to have OS-positivity of all 4-point functions, for all fields, including those of higher spin. Here I only consider scalar 4-point functions. Moreover, OS-positivity implies convergence of conformal OPE on the vacuum because these expansions amount to an orthogonal decomposition of a vector in a Hilbert space.

I pause for some explanatory remarks. In the Euklidean approach the Hilbert space $\mathcal{H}$ is the completion of a space of sequences of test functions f with support in half spaces, and the scalar product is furnished by OS-positivity. $\mathcal{H}$ carries a contractive representation of a sub-semigroup S of

the Euklidean conformal group G^E. S leaves half spaces invariant. Its contractive representations possess an analytic continuation to unitary positive energy representations of G. This was proven in [49], and generalized in [40]; in mathematics a similar method was later introduced by Gel'fand and Gindikin [27]. S possesses a polar decomposition. Such semigroups are nowadays known in mathematics as Ol'shanskii-semigroups [38]. It would be interesting to reformulate axiomatic analyticity as holomorphy on a quadruple of semigroups S which are sewn together along parts of their boundaries.

OS positivity of 4-point functions is the following statement. Write θ for the Euklidean time reflection $x^D \mapsto -x^D$. Consider finite sequences $f = \{f_{i_2 i_1}(\mathbf{x}_2, \mathbf{x}_1)\}$ of test functions which vanish with all their derivatives unless the Euklidean time components satisfy $x_1^D > 0$ and $x_2^D > 0$. Then

$$A_f \equiv \sum_{i_4,i_3,i_2,i_1} \int ... \int d\mathbf{x}_4 ... d\mathbf{x}_1 \qquad (86)$$
$$\bar{f}_{i_3 i_4}(\theta\mathbf{x}_3, \theta\mathbf{x}_4) G_{i_4 ... i_1}(\mathbf{x}_4, ..., \mathbf{x}_1) f_{i_2 i_1}(\mathbf{x}_2, \mathbf{x}_1) \geq 0$$

Let us verify that this follows from OPE (85) of Euklidean 4-point functions, given positivity of the Wightman 2-point functions $\Delta_+^{\chi_k}(p)$ of all the fields ϕ^k in the OPE.

For Euklidean points $x = (x^0, ..., x^{D-1})$, x^0 is imaginary, and Euklidean time reflection θ is equivalent to complex conjugation $x \mapsto \bar{x}$. For Minkowski momenta p, including in particular all $p \in V_+$, the analytic continuation (175) of the $\mathfrak{Q}^u$ kernels has the following reality property.

$$\mathfrak{Q}^u(-p, \tilde{\chi}, \bar{x}_4, \chi_4, \bar{x}_3, \chi_3) = \overline{\mathfrak{Q}^u(p, \tilde{\chi}, x_3, \chi_3, x_4, \chi_4)} \ . \qquad (87)$$

Insert eq. (85) for $G_{i_4 ... i_1}$ in expression (86) for A_f. Define

$$v_k(p) \ = \ \sum_{i_2,i_1} g_k^{i_2 i_1} \int \int d\mathbf{x}_2 \ d\mathbf{x}_1$$
$$f_{i_2 i_1}(\mathbf{x}_2, \mathbf{x}_1) \mathfrak{Q}^u(p, \tilde{\chi}, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1)$$

Then

$$A_f = \sum_k \int_{V_+} (dp) \ \langle \bar{v}_k(p), \Delta_+^{\chi_k}(p) v_k(p) \rangle \geq 0 \ . \qquad (88)$$

Remember that $\langle , \rangle$ stands for the bilinear form which is given by contraction of tensor indices; there is no complex conjugation involved in it.

6 Lessons from Lagrangean field theory

6.1 Renormalized Schwinger Dyson equations

Lagrangean field theories are theories of fundamental fields. For simplicity, assume that there is one such field ϕ^0. In Lagrangean field theory there exists an infinite set of renormalized Schwinger Dyson equations which the collection of all connected n-point Green functions of ϕ^0 must satisfy; it involves no coupling constants [69, 20, 21]. The equations [35] for ϕ^4-theory could be simplified by introducing a second fundamental field $:(\phi^0)^2:$.

Renormalized perturbation theory is the iterative solution of all these equations, together with renormalization conditions which introduce the renormalized coupling constants. When only the equations for the higher n-point functions are solved by iteration, the result is known as a skeleton graph expansions. This leaves the renormalized Schwinger Dyson equations for two- and three-point functions to be solved; they are also known as bootstrap equations in the context of conformal field theory [54, 53, 61, 52].

As an aside: It has been known for some time that gravity has a polynomial action in suitable variables [78]. In the presence of a cosmological term it can be brought to a F^2-form, where F is a de Sitter (or anti de Sitter) field strength [50]. Although they are not in the literature, as far as I know, renormalized Schwinger Dyson equations could be written down, after introducing suitable composites as auxiliary fields, similarly to $:(\phi^0)^2:$ above, assuming that an appropriate gauge fixing can be found. But these equations do not admit an iterative solution. This is in agreement with the conviction that gravity is perturbatively non-renormalizable. The possibility of a conformal invariant short distance behaviour is being discussed [37].

Let us return to conformal field theory. Suppose that the fundamental field ϕ^0 has spin 0 and dimension d_0.

The [totality of] renormalized Schwinger Dyson equations have been solved by Euklidean conformal partial wave expansions [42, 43, 14], and this leads to relations between the square of a coupling constant which appears in a Born term, and the residue of a pole of $g^{(43)(21)}(\chi) = g^{(43)(21)}(\tilde{\chi})$ at $\chi = \chi_0 = [0, d_0]$.

As a consequence, a Born term can be included in the Euklidean partial wave expansion by altering the path of the c-integration as shown in figure 1 a).

When the path of the c-integration is shifted (after a split of the integrand) in the derivation of OPE from the Euklidean partial wave expansion

(section 5.5) there results a cancellation of the contribution of the shadow pole as shown in figure 1 b). This cancellation was first proposed by Ferrara, Gatto, Grillo and Parisi [16], see also [18, 17].

The cancellation is illustrated at the example of ϕ^3-theory in $D = 6 + \epsilon$ dimensions by the result eq.(7) and remarks following it.

6.2 Consequences of Bethe Salpeter equations

Now consider composite fields. Lagrangean field theory teaches that all three point Green functions V with one composite and two fundamental fields should satisfy the Bethe Salpeter equation. In short hand, it reads $BV = V$, where B is the Bethe Salpeter kernel. It has the same form as the Schwinger Dyson equation for the 3-point function in case of a fundamental field.

In conformal theories, V is a sum of kinematically determined three-point functions $\mathfrak{V}^a$ times coupling constants. Currents and stress tensor have been examined in [51] and [15].

Consider fields ϕ_k, with spin and dimension $\chi_k = [l_k, d_k]$. Assuming normalization convention (79), validity of the Bethe Salpeter equation amounts to the requirement $b(\chi_k) = 1$ on the Euklidean partial wave b of the Bethe Salpeter kernel B.

If ϕ^0 is the fundamental field of a Lagrangean field theory, this can be used to give an argument that the physical fields ϕ_k which appear in the OPE of $\phi^0\phi^0\Omega$ are in bijective correspondence with pairs of poles of $g(\chi) \equiv g_{0000}^{(43)(21)}(\chi)$ at $\chi = \chi_k$ and $\chi = \tilde{\chi}_k$.

This result implies that the OPE contains no contributions other than from poles of $g(\chi)$. In particular, the poles from the normalization factor (79) in $N_l(...)N_l(...)[1 + g(\chi)]$ must all be cancelled by zeroes of $[1 + g(\chi)]$.

The argument is based on the Bethe Salpeter equation for the 4-point function. It is an equation for the 4-point Green function G which is 1-particle irreducible in the given channel, i.e. without the above mentioned Born term. It reads, in shorthand, $G = B+BG$. Upon Euklidean partial wave expansion, this translates into $g(\chi) = b(\chi) + b(\chi)g(\chi)$, or $g(\chi) = b(\chi)[1 - b(\chi)]^{-1}$. Therefore, $g(\chi)$ has a pole at $\chi = \chi_k$. Because $g(\chi) = g(\tilde{\chi})$, it also has a pole at $\chi = \tilde{\chi}_k$. .

7 Mellin amplitudes of individual Euklidean partial waves

Here I derive the Mellin representation of an individual Euklidean partial wave I^χ, $\chi = [l, h+c]$, which enters the Euklidean partial wave expansion of the 4-point function of four scalar fields $\phi^{i_1}, ..., \phi^{i_4}$ of dimension $d_1, ..., d_4$. Without loss of generality I assume that the scalar fields are real and that the normalization constants in the 3-point functions are chosen real for real c.

To make contact with the formulae in earlier work, I introduce the notations, for $(ij) = (12), (34)$

$$\begin{aligned}\frac{1}{2}(d_i + d_j) &= h + c_+^{ij} = h + c_+^{ji} \\ \frac{1}{2}(d_i - d_j) &= c_-^{ij} = -c_-^{ji}\end{aligned}$$

By definition

$$\begin{aligned}&I^\chi(d_4, \mathbf{x}_4, ..., d_1, \mathbf{x}_1) = \\ &\int d^D\mathbf{x} \langle \mathfrak{V}(\mathbf{x}, \tilde{\chi}, \mathbf{x}_4, \chi_4, \mathbf{x}_3, \chi_3), \mathfrak{V}(\mathbf{x}, \chi, \mathbf{x}_2, \chi_2, \mathbf{x}_1, \chi_1)\rangle \\ &= (2\pi)^{-D} \int d^D\mathbf{x} N_l(c_+^{34}, c_-^{34}, -c) N_l(c_+^{12}, c_-^{12}, c) \langle f_l^\lambda, f_l^\mu\rangle \\ &\quad (\frac{1}{2}\mathbf{x}_{12}^2)^{-\frac{1}{2}(h-c+l)-c_+^{12}} \ (\frac{1}{2}\mathbf{x}_{34})^{-\frac{1}{2}(h+c+l)-c_+^{34}} \\ &\quad (\frac{1}{2}\mathbf{x}_{10}^2)^{-\frac{1}{2}(h+c-l)-c_-^{12}} \ (\frac{1}{2}\mathbf{x}_{20}^2)^{-\frac{1}{2}(h+c-l)+c_-^{12}} \\ &\quad (\frac{1}{2}\mathbf{x}_{30}^2)^{-\frac{1}{2}(h-c-l)-c_-^{34}} \ (\frac{1}{2}\mathbf{x}_{40}^2)^{-\frac{1}{2}(h-c-l)+c_-^{34}}\end{aligned} \tag{89}$$

with

$$\lambda = 2\left(\frac{\mathbf{x}_{10}}{\mathbf{x}_{10}^2} - \frac{\mathbf{x}_{20}}{\mathbf{x}_{20}^2}\right), \tag{90}$$

$$\mu = 2\left(\frac{\mathbf{x}_{40}}{\mathbf{x}_{40}^2} - \frac{\mathbf{x}_{30}}{\mathbf{x}_{30}^2}\right). \tag{91}$$

f_l^λ is the vector in the representation space V^l of the Euklidean Lorentz group $SO(D)$ associated with the homogeneous polynomial $f_l^\lambda(\mathfrak{z}) = (\lambda\mathfrak{z})^l$.

The bilinear form

$$\langle f_l^\lambda, f_l^\mu \rangle = (\lambda_{\nu_1}...\lambda_{\nu_l} - traces)(\mu_{\nu_1}...\mu_{\nu_l} - traces) \ .$$

f_l^λ is invariant under the $SO(D-1)$ subgroup of rotations which leave λ invariant, and it is homogeneous in λ of degree l. The expression $\langle f_l^\lambda, f_l^\mu \rangle$ is therefore given by the zonal spherical function of $SO(D)$ multiplied with $(|\lambda||\mu|)^l$ times the norm squared $|c_l|^2$ of f_l^e, e= unit vector. It depends on the angle θ between λ and μ, i.e. $\cos\theta = \lambda\mu/|\lambda||\mu|$. The zonal sperical function of a irreducible representation l of $SO(D)$ is the matrix element of the corresponding representation operator $D^l(g)$ for $g \in SO(D)$ between normalized $SO(D-1)$-invariant states. It depends on g only through one angle of rotation θ. It is proportional to the Gegenbauer polynomial C_l^{h-1}. I choose to absorb the normalization factor $|c_l|^2$ into the definition of the zonal spherical function Y_l^D. Explicitly

$$\begin{aligned} \langle f_l^\lambda, f_l^\mu \rangle &= (|\lambda||\mu|)^l Y_l^D(\cos\theta) \qquad (92) \\ &= \frac{l!}{(h-1)_l} 2^{-l} (|\lambda||\mu|)^l C_l^{h-1}(\cos\theta) \\ |c_l|^2 &= 2^{-l} \frac{(2h-2)_l}{(h-1)_l} \qquad (93) \end{aligned}$$

Inserting this result we get

$$\begin{aligned} I^\chi &= (2\pi)^{-D} 2^{3h+c_+^{12}+c_+^{34}-l} N_l(c_+^{34}, c_-^{34}, -c) N_l(c_+^{12}, c_-^{12}, c) \\ &\quad (\mathbf{x}_{12}^2)^{-\frac{1}{2}(h-c)-c_+^{12}} (\mathbf{x}_{34}^2)^{-\frac{1}{2}(h+c)-c_+^{34}} \\ &\quad \int d^D\mathbf{x}_0 (\mathbf{x}_{10}^2)^{-\frac{1}{2}(h+c)-c_-^{12}} (\mathbf{x}_{20}^2)^{-\frac{1}{2}(h+c)+c_-^{12}} \\ &\quad\quad (\mathbf{x}_{30}^2)^{-\frac{1}{2}(h-c)-c_-^{34}} (\mathbf{x}_{40}^2)^{-\frac{1}{2}(h-c)+c_-^{34}} Y_l^D(\cos\theta) \ . \end{aligned}$$

The integral can be evaluated with the help of the generalized Symanzik 4-star formula (162) of section 12.1 with

$$\begin{aligned} \delta_1 &= \frac{1}{2}(h+c) + c_-^{12} \ , \\ \delta_2 &= \frac{1}{2}(h+c) - c_-^{12} \ , \\ \delta_3 &= \frac{1}{2}(h-c) + c_-^{34} \ , \\ \delta_4 &= \frac{1}{2}(h-c) - c_-^{34} \ , \qquad (94) \end{aligned}$$

The result involves variables which I denote by δ'_{ij} subject to the constraints $\sum_j \delta'_{ij} = \delta_i$, and an integration over imaginary parts s_1, s_2 of two independent ones among the so constrained variables. I^χ comes out to equal

$$\pi^{-h} 2^{c_+^{12}+c_+^{34}-l+h} N_l(c_+^{34}, c_-^{34}, -c) N_l(c_+^{12}, c_-^{12}, +c)$$
$$(2\pi i)^{-2} \int_{-i\infty}^{i\infty} \int_{-i\infty}^{i\infty} ds_1 ds_2 \mathfrak{P}_l^D(\{\delta'_{ij}\}) \prod{}' \Gamma(\delta'_{ij}) (\mathbf{x}_{ij}^2)^{-\delta'_{ij}}$$
$$\Gamma(\delta'_{12} - \frac{1}{2}l)(\mathbf{x}_{12}^2)^{-\delta'_{12}-\frac{1}{2}(h-c)-c_+^{12}}$$
$$\Gamma(\delta'_{34} - \frac{1}{2}l)(\mathbf{x}_{34}^2)^{-\delta'_{34}-\frac{1}{2}(h+c)-c_+^{34}}$$

The product $\prod'$ runs over the four links $(13), (14), (23), (24)$.

Introduce new variables

$$\begin{aligned} \delta_{12} &= \delta'_{12} + \frac{1}{2}(h+c) + c_+^{12} \\ \delta_{34} &= \delta'_{34} + \frac{1}{2}(h-c) + c_+^{34} \\ \delta_{ij} &= \delta'_{ij}, \qquad (ij) \neq (12), (34) \ . \end{aligned} \tag{95}$$

They satisfy the constraints $\sum_j \delta_{ij} = d_i$.

We may compare with the general form (47) of the Mellin representation of a 4-point function to deduce the Mellin amplitude $\mathfrak{M}_\chi$ of a Euklidean partial wave I^χ,

$$\begin{aligned} &\mathfrak{M}_\chi(\{\delta_{ij}\}) = N_l(c_+^{34}, c_-^{34}, -c) N_l(c_+^{12}, c_-^{12}, c) \tilde{\mathfrak{M}}_\chi(\{\delta_{ij}\}) \ , \\ &\tilde{\mathfrak{M}}_\chi(\{\delta_{ij}\}) = (\pi/2)^{-h} 2^{-l} \mathfrak{P}_l^D(\{\delta'_{ij}\}) \\ &\frac{\Gamma(\delta_{12} - \frac{1}{2}(h+c+l) - c_+^{12})}{\Gamma(\delta_{12})} \frac{\Gamma(\delta_{34} - \frac{1}{2}(h-c+l) - c_+^{34})}{\Gamma(\delta_{34})} \end{aligned} \tag{96}$$

where δ'_{ij} is supposed to be expressed in terms of δ_{ij} using eqs.(95)ff, and the polynomials $\mathfrak{P}_l^D$ are defined in Appendix 12.

Let us express the result in terms of the independent variables β_{12} and $\gamma_{12} = \beta_{23} - \beta_{13}$ which determine the variables δ_{ij} according to the rules of section 3.2, given the field dimensions $d_1, ..., d_4$. It is helpful to use, in place of β_{12} the shifted variable

$$\delta^{(43)(21)} = \beta_{12} - \frac{1}{6}(d_1 + d_2 + d_3 + d_4) = -\frac{1}{2} s_{12} \ .$$

It is related to the Mandelstam variable s_{12} of the introduction. Its significance transpires from the relations

$$\delta_{12} - \frac{1}{2}(d_1 + d_2) = \delta^{(43)(21)} = \delta_{34} - \frac{1}{2}(d_3 + d_4) \tag{97}$$

One computes

$$\begin{aligned}
\delta'_{12} &= \delta^{(43)(12)} + \frac{1}{2}(h + c) \ , \\
\delta'_{34} &= \delta^{(43)(12)} + \frac{1}{2}(h - c) \ , \\
\delta'_{13} &= -\frac{1}{2}\delta^{(43)(12)} + \frac{1}{2}(c_-^{12} + c_-^{34} - \gamma_{12}) \ , \\
\delta'_{14} &= -\frac{1}{2}\delta^{(43)(12)} + \frac{1}{2}(c_-^{12} - c_-^{34} + \gamma_{12}) \ , \\
\delta'_{23} &= -\frac{1}{2}\delta^{(43)(12)} + \frac{1}{2}(-c_-^{12} + c_-^{34} + \gamma_{12}) \ , \\
\delta'_{24} &= -\frac{1}{2}\delta^{(43)(12)} + \frac{1}{2}(-c_-^{12} - c_-^{34} - \gamma_{12}) \ ,
\end{aligned} \tag{98}$$

Define, for $\chi = [l, h + c]$ the polynomial in γ_{12} which depends parametrically on the differences of dimensions $c_-^{12} = \frac{1}{2}(d_1 - d_2)$ and $c_-^{34} = \frac{1}{2}(d_3 - d_4)$, and on $\delta^{(43)(21)}$ by

$$\mathfrak{P}^D_\chi(c_-^{12}, c_-^{34}; \delta^{(43)(21)}|\gamma_{12}) = \mathfrak{P}^D_l(\{\delta'_{ij}\}) \ , \tag{99}$$

evaluated at $\{\delta'_{ij}\}$ as given by eq.(98). Then

$$\begin{aligned}
&\tilde{\mathfrak{M}}_\chi(\{\delta_{ij}\}) = (\pi/2)^{-h} 2^{-l} \mathfrak{P}^D_\chi(c_-^{12}, c_-^{34}; \delta^{(43)(21)}|\gamma_{12}) \\
&\frac{\Gamma(\delta^{(43)(21)} + \frac{1}{2}(h - l) - \frac{1}{2}c)}{\Gamma(\delta_{12})} \frac{\Gamma(\delta^{(43)(21)} + \frac{1}{2}(h - l) + \frac{1}{2}c)}{\Gamma(\delta_{34})} \ .
\end{aligned} \tag{100}$$

We see that $\tilde{\mathfrak{M}}_\chi$ has two families of poles in δ_{12} which are related to each other by the substitution $c \mapsto -c$, and it is a polynomial of degree l in the other independent variable γ_{12}.

The factors $\Gamma(\delta_{12})^{-1}\Gamma(\delta_{34})^{-1}$ were left in this form in order not to obscure the cancellation which occurs in the Mellin representation (47), and which ensure that the reduced Mellin amplitude $\hat{M}^c$, which is related to M^c by eq. (46), does not have extra poles at non-positive integer δ_{12} and δ_{34}.

After splitting the pairs of poles and shifting the path of the c-integration as described in section 8, expression (100) will (only) produce the poles in M^c

coming from fields of Lorentz spin l in the OPE. It is nevertheless amusing to compare with the structure of the Veneziano Beta-function $B(-\alpha(s),-\alpha(t))$. Its poles in s come from a quotient of Γ-functions $\Gamma(-\alpha(s))/\Gamma(-\alpha(s)-\alpha(t))$, so that their positions are obtained by a shift from the positions of its zeroes.

It follows from the amputation identity (71) that

$$\mathfrak{M}_\chi = \mathfrak{M}_{\tilde{\chi}} \ .$$

This reflects the equivalence of representations χ and $\tilde{\chi}$. This implies the same for $\tilde{\mathfrak{M}}_\chi$ up to a factor. One finds by inspection that

$$\mathfrak{P}^D_\chi(c_-^{12},c_-^{34};\delta^{(43)(21)}|\gamma) = \mathfrak{P}^D_{\tilde{\chi}}(c_-^{34},c_-^{12};\delta^{(43)(21)}|\gamma) \ . \tag{101}$$

Therefore

$$\tilde{\mathfrak{M}}_\chi = \tilde{\mathfrak{M}}_{\tilde{\chi}} \ \text{ if } c_-^{12} = c_-^{34}. \tag{102}$$

To verify eq.(101) note that $c \leftrightarrow -c$, $c_-^{12} \leftrightarrow c_-^{34}$ takes $\delta_1 \leftrightarrow \delta_3$, $\delta_2 \leftrightarrow \delta_4$ and $\delta'_{34} \leftrightarrow \delta'_{12}$. The result now follows from eqs.(159,160) by a change of summation variables $k_{14} \leftrightarrow k_{23}$.

The coefficient of the leading power in γ in the polynomial $\mathfrak{P}^D_\chi(c_-^{12},c_-^{34};\delta^{(43)(21)}|\gamma)$ can be evaluated in closed form, starting from the defining equations (159) and (160) and using expressions (98) for δ'_{ij}.To leading order in γ, $\prod'(\delta_{ij})_{k_{ij}} = (-)^{k_{13}+k_{24}}\prod'(\gamma/2)^{k_{ij}}$. Inserting the standard integral representation (8.381) of [30], the summations over the four k_{ij} can be performed with the help of the multinomial theorem. The result reads

$$\begin{aligned}&\mathfrak{P}^D_\chi(c_-^{12},c_-^{34};\delta^{(43)(21)}|\gamma) = \\ &a_0^l(D)(h+c-1)_l(h-c-1)_l\mathfrak{p}^D_\chi(c_-^{12},c_-^{34})(-\gamma/4)^l + ...\end{aligned} \tag{103}$$

with

$$\begin{aligned}\mathfrak{p}^D_\chi(c_-^{12},c_-^{34}) \ &= \ \Gamma(\frac{1}{2}(h+c)+c_-^{12})^{-1}\Gamma(\frac{1}{2}(h+c)-c_-^{12})^{-1} \\ &\quad \Gamma(\frac{1}{2}(h-c)+c_-^{34})^{-1}\Gamma(\frac{1}{2}(h-c)-c_-^{34})^{-1}\end{aligned}$$

The important observation is that the dependence on c_-^{12} and on c_-^{34} in $\mathfrak{p}(c_-^{12},c_-^{34})$ factorizes.

The result (103)f is in agreement with the relation (101). It permits to extract the constant factor which relates $\mathfrak{P}^D_\chi(c_-^{12},c_-^{34};\delta^{(43)(21)}|\gamma)$ and $\mathfrak{P}^D_{\tilde{\chi}}(c_-^{12},c_-^{34};\delta^{(43)(21)}|\gamma)$. The same factor is obtained from eq.(96) and expressions in the literature [11] for the normalization factors.

In later applications of the result (103), χ_k will substitute for χ; it will be given by spin and dimension of some field in the OPE.

8 Poles in the Mellin amplitude

8.1 Dynamical poles from poles in the partial wave

Inserting the Mellin representation (100) of the individual partial waves into the partial wave expansion (73), one finds that the connected 4-point Green function admits a Mellin representation with Mellin amplitude

$$M^c(\{\delta_{ij}\}) = \int d\chi \; g^{(43)(21)}(\chi) \; \mathfrak{M}_\chi(\{\delta_{ij}\}). \tag{104}$$

$\chi = [l, h + c]$, and this formula involves a sum over l and an integration over c along the imaginary axis (or along another path if there is a Born term, see below).

OPE imply asymptotic expansions of the full Greens functions, *not* of their connected parts. Nonvanishing disconnected parts $G_{i_4 i_2}(\mathbf{x}_4, \mathbf{x}_2) G_{i_3 i_1}(\mathbf{x}_3, \mathbf{x}_1)$ and $G_{i_4 i_1}(\mathbf{x}_4, \mathbf{x}_1) G_{i_3 i_2}(\mathbf{x}_3, \mathbf{x}_2)$ require that $d_4 = d_2, d_3 = d_1$ and $d_4 = d_1, d_3 = d_2$, respectively. We discuss first the case when neither of these two disconnected parts is present.

Both the amplitude $g^{(43)(21)}(\chi)$ and the normalization factors N_l depend on normalization conventions, but the product $N_l(c_+^{34}, c_-^{34}, -c) N_l(c_+^{12}, c_-^{12}, c) g^{(43)(21)}(\chi)$ does not. We know from section 5.5 that non-derivative Lorentz-irreducible fields ϕ^k with spin and dimension $\chi_k = [l_k, d_k = h + c_k]$ which appear in the OPE of both $\phi^{i_1} \phi^{i_2}$ and of $\phi^{i_3 *} \phi^{i_4 *}$ are in bijective correspondence with pairs of poles in $N_l(c_+^{34}, c_-^{34}, c) N_l(c_+^{12}, c_-^{12}, -c) \; g^{(43)(21)}(\chi)$ for $l = l_k$ in c at $c = \pm c_k$. A complete derivation of the result was published in ref.[14] for even space time dimension D, and under some restrictions on the field dimensions. It was stated that the result can be generalized.

We wish to avoid restrictions on differences c_-^{12} and c_-^{34} of dimensions of the external fields, because factorization of 4-point functions is a constraint on the dependence of residues on the external fields. For the canonical choice of normalization factors [11] which is used in ref. [14], the product of normalization factors $N_l(c_+^{34}, c_-^{34}, -c) N_l(c_+^{12}, c_-^{12}, c)$ has branch cuts, for general c_-^{12}, c_-^{34}. Therefore it is only $\tilde{\mathfrak{M}}_\chi$ rather than $\mathfrak{M}_\chi$ which is meromorphic. We need to extract the product of normalization factors from $\mathfrak{M}_\chi$. It is convenient to postpone this, however.

Our object is to show that to each of the aforementioned pairs of poles there corresponds a family of poles of $M^c(\{\delta_{ij}\})$ in the variable β_{12}, or equiv-

alently, in $\delta^{(43)(21)} = -\frac{1}{2}s_{12} = \delta_{12} - \frac{1}{2}(d_1 + d_2)$ at positions

$$\begin{aligned} \delta^{(43)(21)} &= \delta_l^n(d_k) \ , \qquad n = 0, 1, 2, ... \\ \delta_l^n(d) &= -\frac{1}{2}(d - l) - n \ . \end{aligned} \tag{105}$$

Poles of M^c in the β_{ij} can only arise when the path of the c-integration in the integral (104) gets pinched between singularities of the integrand when β_{ij} is moved in the complex plane. It follows then from the polynomial character of the factor $\mathfrak{P}_\chi^D$ in the expression (100) for $\tilde{\mathfrak{M}}_\chi$ that the only poles are in β_{12} and their residues are polynomials of degree l in the remaining independent variable $\gamma_{12} = \beta_{23} - \beta_{13}$.

To get more explicit results, the symmetry in $c \mapsto -c$ is exploited in the same way as in the derivation of OPE from Euklidean partial wave expansions. The first step was to decompose the Euklidean partial wave into two parts such that the second part is obtained from the first by substitution $c \mapsto -c$, and the strength of the singularity of the first part at $x_{12}^2 = 0$ decreases as the real part of c is increased; this is achieved by the split (81) of the 3-point function. It is analog to the split of Legendre functions $P_{-\frac{1}{2}+i\sigma}$ of the first kind into Legendre functions of the second kind which appears in the group theoretical approach to Regge theory [64]. The corresponding decomposition of the Mellin representation $\mathfrak{M}_\chi$ of the Euklidean partial wave reads

$$\begin{aligned} \mathfrak{M}_\chi &= \frac{\pi}{\sin \pi(c + l)} \big[N_l(c_+^{34}, c_-^{34}, -c) N_l(c_+^{12}, c_-^{12}, c) \mathfrak{M}_\chi^+ \\ &\quad - N_l(c_+^{34}, c_-^{34}, c) N_l(c_+^{12}, c_-^{12}, -c) \mathfrak{M}_{\tilde{\chi}}^+ \big] \end{aligned} \tag{106}$$

with the requirement on $\mathfrak{M}_\chi^+$ that the poles in c of $\mathfrak{M}_\chi^+$ occur only at the positions of the poles of $\Gamma(\delta_{12} - \frac{1}{2}(h - c + l) - c_+^{12})$, i.e. when the condition (105) is fulfilled. For the canonical choice [11], the ratio of the two products of normalization factors in eq.(106) is meromorphic; therefore $\mathfrak{M}_\chi^+$ shares the meromorphy properties of $\tilde{\mathfrak{M}}_\chi$.

Since we know that poles in β_{12} arise only from pinches, it suffices to represent $\tilde{\mathfrak{M}}_\chi$ as a formal sum of pole terms in c labeled by $n = 0, 1, 2, ...$ [5]

[5] This is true in spite of the fact that the formal sum of pole terms does not indicate the correct asymptotic behavior as the imaginary part of c tends to ∞. We know from the work in [14] that the path of the c-integration in the Euklidean partial wave expansion can be shifted, after the split. This results from the good asymptotic behavior of the $\mathfrak{Q}$-kernels

Using that $\Gamma(-n-c) = \pi[\sin\pi(-c-n)\Gamma(1+n+c)]^{-1}$, we find from the requirements on its poles that

$$\begin{aligned}\Gamma(\delta_{12})\Gamma(\delta_{34})\mathfrak{M}^+_\chi &= -(\pi/2)^{-h}\sum_{n=0}^{\infty}\frac{(-2)^{-l}}{n!\Gamma(n+c+1)}\\ &(\delta^{(43)(21)} - \delta_l^n(h+c))^{-1}\mathfrak{P}^D_\chi(c^{12}_-, c^{34}_-; \delta_l^n(h+c)|\gamma_{12})\\ &+\text{nonsingular}\end{aligned} \tag{107}$$

Remember that the factor $\Gamma(\delta_{12})\Gamma(\delta_{34})$ cancels in the Mellin representation (47).

Inserting the split (106) and making a change of variables $c \mapsto -c$ in the second term, the integral (104) becomes

$$\begin{aligned}M^c(\{\delta_{ij}\}) &= \frac{2\pi}{\sin\pi(c+l)}\int d\chi\\ &N_l(c^{34}_+, c^{34}_-, -c)N_l(c^{12}_+, c^{12}_-, c)\; g^{(43)(21)}(\chi)\mathfrak{M}^+_\chi(\{\delta_{ij}\})\;.\end{aligned}$$

Using the pole structure of $\mathfrak{M}^+$ as given by eq.(107) we can identify the pinches which lead to poles in $\delta^{(43)(21)}$.

Let us initially ignore the poles of $\sin\pi(c+l)$. It will turn out that they make no contribution.

Consider fixed n and start with $\delta^{(43)(21)}$ large and positive. Then the pole of $(\delta^{(43)(21)} + \frac{1}{2}(h+c-l)+n)^{-1}$ in c is at large negative c. As we move $\delta^{(43)(21)}$ in negative direction, the pole will eventually touch the path of c-integration. We can deform the path of c-integration to avoid the pole, until the path gets pinched between the pole of $(\delta^{(43)(12)} + \frac{1}{2}(h+c-l)+n)^{-1}$ and of $N_l(c^{34}_+, c_-, c)N_l(c^{12}_+, c_-, -c)\; g^{(43)(21)}([l, h+c])$ at $c = c_k$.

Suppose now that the residue of the latter pole factorizes as in eq.(84)

The pinch will lead to a pole in M^c at position $\delta^{(43)(21)} = \delta_l^n(h+c_k)$ with residue given by

$$\begin{aligned}\Gamma(\delta_{12})\Gamma(\delta_{34})res\; M^c &= \bar{g}_k^{i_3 i_4} g_k^{i_1 i_2} \qquad (108)\\ &\quad \mathfrak{r}^n_{\chi_k}\mathfrak{P}^D_{\chi_k}(c^{12}_-, c^{34}_-; \delta_l^n(h+c_k)|\gamma_{12}).\\ \mathfrak{r}^n_\chi &= -(\pi/2)^{-h}\frac{(-2)^{-l}}{n!\Gamma(n+c+1)} \qquad (109)\\ \chi_k &= [l, h+c_k]\;.\end{aligned}$$

If the OPE contains fields ϕ^a, ϕ^b of the same Lorentz spin l which differ in dimension by an even integer $d_b - d_a = 2N$, then a coincidence of the positions of poles $\delta_l^n(d_a) = \delta_l^{(n+N)}(d_b)$ occurs.

It remains to discuss possible contributions from pinches with poles of $1/\sin\pi(c+l)$ at integer c. The same problem occurs in the derivation of OPE from Euklidean partial wave expansions and was discussed in ref.[14], for even D. The result is that all such contributions cancel in a subtle way, partly because of zeroes of the Plancherel measure and partly because of partial equivalences of elementary representations χ at so-called exceptional integer points. For odd D, the cancellation follows from zeroes of the Plancherel weight ρ_l at integer c. But in this case the Plancherel weight has zeroes at half odd integral values of c, whose contributions need to be cancelled either by zeroes of $g^{(43)(21)}(\chi)$, or by partial equivalences among exceptional integer points. [Discrete series representations cannot occur for scalar amplitudes].

Let us now turn to situations with disconnected parts. Consider the extreme case of four equal real fields ϕ of dimension $d_4 = d_3 = d_2 = d_1 = h + c_+$ where all three disconnected parts are present. One of them represents the contribution of the unit operator to OPE. We have now $c_+^{34} = c_+^{12} = c_+$ and $c_- = 0$. We assume that normalization conventions are so chosen that the Euklidean partial wave expansion of the 4-point function involves

$$N_l(c_+, 0, c)N_l(c_+, 0, -c)[1 + g^{(43)(21)}([l, h + c])] \ , \tag{110}$$

where 1 is the contribution of the disconnected part. The appropriate product of normalization factors is given in eq.(79); it is a meromorphic function of c.

In this situation, the fields in the OPE of $\phi\phi$ with Lorentz spin l and dimension $d_k = h + c_k$ are in bijective correspondence with pairs of poles of expression (110) at $c = \pm c_k$. We distinguish two sets of poles of
$N_l(c_+, 0, c)N_l(c_+, 0, -c)g^{(43)(21)}([l, h + c])$.

i) poles of
$N_l(c_+, 0, c)N_l(c_+, 0, -c)[1 + g^{(43)(21)}([l, h + c])]$
I call these dynamical poles

ii) poles of $N_l(c_+, 0, c)N_l(c_+, 0, -c)$ at which $g([l, h + c]) = -1$.
I call these kinematical poles.

The dynamical poles make contributions as discussed before; in the expression for the residue, $[1 + g^{(43)(21)}([l, h + c])]$ should be substituted for $g^{(43)(21)}([l, h + c])$.

8.2 Kinematical poles in M^c

In the generic case of anomalous dimensions, when the disconnected parts have asymptotic expansion which involve integer powers of x_{12}^2 which are incompatible with OPE, the purpose of the kinematical poles is to cancel the disconnected parts of the full amplitude. As will be shown right below, it follows from this that it must be true that $g(\chi) = -1$ at all the poles of $N_l(c_+, 0, c)N_l(c_+, 0, -c)$. Therefore, all the poles of $N_l(c_+, 0, c)N_l(c_+, 0, -c)[1+ g^{(43)(21)}([l, h + c])]$ will actually be poles of $[1 + g(\chi)]$. Another argument for this, which is valid when ϕ is the fundamental field of a Lagrangean field theory, was given in section 6.2.

To see the cancellation, compare with the generalized free field theory which has the same disconnected parts as the interacting theory, but zero connected part; its conformal partial wave expansion is given by eq.(75) with $g(\chi) \equiv 0$. Inserting the Mellin representation $\mathfrak{M}_\chi$ of individual partial waves, we end up with a sum over l and an integral over c. To avoid divergences, we imagine that the summation over l is done at the end. Repeating the procedure above, we get poles in the Mellin representation of the contributions from individual l from the poles of $N_l(c_+, 0, c)N_l(c_+, 0, -c)$ which are the negative of the contributions from the contribution of kinematical poles to the connected amplitude M^c in the interacting theory.

Because positivity is a property of the full Wightman functions, it follows from this cancellation that the contributions from the kinematical poles to the connected amplitude do not violate positivity.

8.3 Born terms

A word should be added concerning the case, when a fundamental field ϕ^f of dimension $h + c_f$, $c_f < 0$ contributes to the OPE. This happens in ϕ^4-theory in 3 dimensions, for instance, with $\phi^f =: \phi^2 :$. As discussed in section 6.1 the initial path of the c-integration is different in this case, see figure 1 a), because it is possible to include the Born term by altering the path of the c-integration.

Because of the symmetry of the partial wave $g^{(43)(21)}([l, h + c])$ under $c \mapsto -c$, the situation is unchanged after the split (106) and subsequent substitution $c \mapsto -c$ in the second term. Upon shifting the path of the c-integration, there will be a contribution from the pole at $c = c^f$ which reproduces the contribution of ϕ^f to the OPE, but there will be no contribution

from its shadow at $c = -c^f$, see figure 1 b).

9 Factorization of scalar 4-point functions

In the following I consider the Mellin amplitudes of 4-point functions of scalar fields $\phi^{i_4}, ..., \phi^{i_1}$. They may be regarded as functions of two independent variables β_{12} and $\gamma_{12} = \beta_{23} - \beta_{13}$. Fields ϕ_k which appear in the OPE of $\phi^{i_2}\phi^{i_1}$ and $\phi^{i_3 *}\phi^{i_4 *}$ are symmetric tensor fields of some rank l_k and some dimension d_k. As we saw, they determine integer spaced families of poles in β_{12} at positions of $\beta_{12} = \delta^{(43)(21)} + \frac{1}{6}(d_1 + d_2 + d_3 + d_4)$ given by $\delta^{(43)(21)} = \delta^n_{l_k}(h + c_k) = -\frac{1}{2}(h + c_k - l_k) - n$. Their residues are polynomials of degree l_k in γ_{12}. Factorization is a statement about the $i_4, ..., i_1$ dependence of these residues which follows from OPE.

First I explain how the spin and dimension $\chi_k = [l_k, d_k]$ of all the fields is read off from the poles of $M^c_{i_4 ... i_1}$. In doing so, the amplitude $M^c_{i_4 ... i_1}$, regarded as formal sums of pole terms, will be rewritten so that the integer spaced families of poles which correspond to individual fields ϕ_k will be exhibited, as in eq.(111) below. I write $c^{12}_- = d_{i_1} - d_{i_2}$ etc. for differences of dimensions of the external fields.

$$\begin{aligned} M^c_{i_4 ... i_1} &= \Gamma(\delta_{12})^{-1}\Gamma(\delta_{34})^{-1} \sum_k \sum_{n=0}^{\infty} g_k^{i_4 ... i_1} \mathfrak{r}^n_k \\ &\quad \mathfrak{P}^D_{\chi_k}(c^{12}_-, c^{34}_-; \delta^n_{l_k} | \gamma_{12}) \frac{1}{\delta^{(43)(21)} - \delta^n_{l_k}(d_k) - n} \end{aligned} \tag{111}$$

Herein, $\mathfrak{P}$ are the kinematically determined polynomials in γ_{12} defined by eq.(99), using the Appendix (section 12), and $\mathfrak{r}$ are kinematically determined factors which govern the n-dependence of the residues of satellite poles $n = 1, 2, ...$, see eq.(109). There is a dependence on differences of external dimensions in these quantities.

I assume that the normalization of fields is so chosen that the normalization factor $\mathbf{c}_k$ of their two-point function is $+1$ for physical fields, and -1 for ghosts.

Factorization now requires that the coefficients $g_k^{i_4 ... i_1}$ factorize in products

$$g_k^{i_4 ... i_1} = \bar{g}_k^{i_4 i_3} g_k^{i_1 i_2} \tag{112}$$

for physical fields ϕ^k. In the case of a ghost, there would be an overall $-$-sign.

Let me explain how the spin and dimension $\chi_k = [l_k, d_k]$ of the fields and the coefficients $g_k^{i_4 \ldots i_1}$ are determined from the poles of $M^c_{i_4 \ldots i_1}$.

I assume again that field dimensions are anomalous, so that there is at most a finite number of fields with given twist $d_k - l_k$. Nevertheless there can be limit points of poles, as mentioned in the introduction. The procedure outlined below must then be carried out in an appropriate order, as specified below, in order to find the fields in all trajectories associated with the different limit points.

Consider what remains of $M^c_{i_4 \ldots i_1}$, considered as a sum of pole terms, after the contributions of zero or some fields ϕ^j in the OPE are subtracted. These contributions have the form of individual terms in the sum over k in the right hand side of eq.(111). Suppose that the remainder has a leading pole in β_{12} (i.e. the pole with largest β_{12}) whose residue is a polynomial in γ_{12} of degree l_k, at a position which is such that $\delta^{(43)(21)} = \delta^0_{l_k}(d_k) \equiv -\frac{1}{2}(d_k - l_k)$; this determines d_k. Thereby spin l_k and dimension d_k of a field ϕ_k are determined which has to be introduced in order to reproduce this pole. Abbreviate $\chi_k = [l_k, d_k]$ as usual. The coefficient of the highest power $-(\gamma_{12}/4)^{l_k}$ of the polynomial residue can be written in the form

$$g_k^{i_4 \ldots i_1} \mathfrak{r}_{\chi_k} a_0^{l_k}(D)(d_k - 1)_{l_k}(D - d_k - 1)_{l_k} \mathfrak{p}^D_{\chi_k}(c_-^{12}, c_-^{34})$$

with some suitable coefficient $g_k^{i_4 \ldots i_1}$ which depends on the labels $i_4 \ldots i_1$ of the external fields.[The quantities a_0^l are the coefficients in the power series expansion (157) of the zonal spherical function, while $\mathfrak{r}$ and $\mathfrak{p}$ are kinematically determined quantities defined in eqs. (109), and (103)f.] Now subtract the contribution of this field in the OPE from the remainder of $M^c_{i_4 \ldots i_1}$. It follows from the definition (103) of $\mathfrak{p}^D_{\chi_k}(c_-^{12}, c_-^{34})$ as coefficient of the leading power $(-\gamma_{12}/4)^{l_k}$ in the polynomial $\mathfrak{P}_{\chi_k}(c_-^{12}, c_-^{34}; \delta^{(43)(21)}|\gamma_{12})$ that the leading pole of what remains afterwards will either have a smaller position β_{12} or a residue of lower order.

If there are no limit points of poles, all poles can be subtracted recursively, typically in infinitely many steps. In the case with limit points, one must proceed in a slightly different order, removing successively the finitely many poles outside smaller and smaller neighborhoods of the limit points and of $\beta_{12} = \infty$.

10 Asymptotic expansions from the Mellin representation

Like the Mellin representation itself, the formulas of this section are valid for arguments throughout the axiomatic domain of analyticity (the permuted extended tube.) If x_i are in Minkowski space, appropriate $i\epsilon$-prescriptions are to be imposed to obtain the Wightman function with a particular ordering of the fields. If x_i are Euklidean, $-x_{ij}^2 = \mathbf{x}_{ij}^2$.

Here I examine the asymptotic expansion of a 4-point function with Mellin representation (47) as $x_{12} \mapsto 0$. It is determined by the poles of M^c in δ_{12}.

Remember that the exponents δ_{ij} in the Mellin representation of a scalar 4-point-function have the form $\delta_{ij} = \delta_{ij}^0 + \beta_{ij}$, where δ_{ij}^0 are kinematically determined by the dimensions d_i, $i = 1, \dots d_4$ of the fields whose 4-point function is considered, and β_{ij} are integration variables, two of which are independent. As independent variables we may take β_{12} and γ_{12},

$$\gamma_{12} = \beta_{23} - \beta_{13} \tag{113}$$

The exponents β_{ij} are expressed through the independent ones as

$$\begin{aligned} \beta_{13} &= -\frac{1}{2}\beta_{12} - \frac{1}{2}\gamma_{12} &&= \beta_{24} \\ \beta_{23} &= -\frac{1}{2}\beta_{12} + \frac{1}{2}\gamma_{12} &&= \beta_{14} \end{aligned} \tag{114}$$

The measure in the Mellin representation is $d\delta \equiv d^2\beta = \frac{1}{2} d\beta_{12} d\gamma_{12}$.

We examine here the particular channel (43)(21), and for this purpose it is again convenient to use in place of β_{12} the shifted variable $\delta^{(43)(21)} = \beta_{12} - \frac{1}{6}(d_1 + d_2 + d_3 + d_4)$ $(= -\frac{1}{2}s_{12})$ so that

$$\delta_{12} - \frac{1}{2}(d_1 + d_2) = \delta^{(43)(21)} = \delta_{34} - \frac{1}{2}(d_3 + d_4)\ . \tag{115}$$

Suppose the Mellin amplitude $M_a^c(\{\delta_{ij}\})$ of a connected scalar 4-point function G_a^c has a pole in β_{12} at position $\delta^{(43)(21)} = \hat{\delta}$ with residue $P(\hat{\delta}|\gamma_{12})$ which is a polynomial in γ_{12}. In applications it will depend on the position $\hat{\delta}$ of the pole.

When the path of the β_{12}-integration in the Mellin representation is shifted past such a pole, a contribution from the residue is obtained which is

equal to

$$R(\hat{\delta}|x_4,\ldots,x_1) = \Gamma(\hat{\delta}_{12})\left(-\frac{1}{2}x_{12}^2\right)^{-\hat{\delta}_{12}} \Gamma(\hat{\delta}_{34})\left(-\frac{1}{2}x_{34}^2\right)^{-\hat{\delta}_{34}} (-2\pi i)^{-1}\int_{-i\infty}^{i\infty}\frac{1}{2}d\gamma_{12}P(\hat{\delta}|\gamma_{12})\underline{\overline{X}} \tag{116}$$

with

$$\underline{\overline{X}} = \prod_{(ij)}{}'\Gamma(\delta_{ij})\left(-\frac{1}{2}x_{ij}^2\right)^{-\delta_{ij}} \tag{117}$$

Herein, $\hat{\delta}_{12}$ and $\hat{\delta}_{34}$ are determined by $\hat{\delta} = \delta^{(43)(21)}$ through eq.(115), and the product $\prod'$ runs over the four unordered pairs $(13),(23),(14),(24)$, i.e. all except (12) and (34). In eq.(117), $\delta_{ij} = \delta_{ij}^0 + \beta_{ij}$ with β_{ij} as in eq.(114).

The factors in the first line may become singular when $x_{12} \mapsto 0$ and/or $x_{34} \mapsto 0$, while the factor $\underline{\overline{X}}$ is not singular when x_1, x_2 stay away from x_3, x_4.

It remains to perform the γ_{12}-integration.

The following well known identity can be used to combine the product of four x-dependent factors into a single factor, at the cost of introducing some variables of integration.

$$\Gamma(\nu)A^{-\nu}\Gamma(\mu)B^{-\mu} = \int_0^1 duu^{\nu-1}(1-u)^{\mu-1}\Gamma(\mu+\nu) (uA+(1-u)B)^{-\nu-\mu} \ . \tag{118}$$

We combine first the x_{13}-dependent factor with the x_{14}-dependent factor, using integration variable u_1, and the remaining two factors using integration variable u_2. As a result, there appears a product of two propagators,

$$\underline{\overline{X}} = \int\int du_1du_2u_1^{\delta_{13}-1}(1-u_1)^{\delta_{14}-1}u_2^{\delta_{24}-1}(1-u_2)^{\delta_{23}-1} \Gamma(\delta_{13}+\delta_{14})\left(-\frac{1}{2}u_1x_{13}^2-\frac{1}{2}(1-u_1)x_{14}^2\right)^{-\delta_{13}-\delta_{14}} \Gamma(\delta_{23}+\delta_{24})\left(-\frac{1}{2}u_2x_{24}^2-\frac{1}{2}(1-u_2)x_{23}^2\right)^{-\delta_{23}-\delta_{24}} . \tag{119}$$

Let us introduce

$$x_5(u) = ux_3 + (1-u)x_4 \tag{120}$$

As u varies from 0 to 1, the point x_5 runs along the straight line from x_3 to x_4. Defining $x_{15}(u) = x_1 - x_5(u)$ and $x_{25}(u) = x_2 - x_5(u)$, one can rewrite the arguments of the above two propagators by using identities like

$$ux_{i3}^2 + (1-u)x_{i4}^2 = x_{i5}(u)^2 + u(1-u)x_{34}^2 \tag{121}$$

for $i = 1, 2$. This shows that expression (119) is a product of two factors of the form $\Gamma(\nu)\left(-\frac{1}{2}x_{i5}^2 - \frac{1}{2}m_{i5}^2\right)^{-\nu}$, which we can interpret as propagators from x_1 to $x_5(u_1)$ and from x_2 to $x_5(1-u_2)$. These "propagators" do not have good positivity properties, though.

In the special case $u_1 = 1 - u_2$ the two points $x_5(u_1)$ and $x_5(1-u_2)$ coincide, and another application of identity (118) (with integration variable denoted v) would produce a formula with a single propagator connecting

$$x_0(v) = vx_1 + (1-v)x_2 \tag{122}$$

to $x_5(u_1)$. The point $x_0(v)$ runs along the straight line connecting x_1 and x_2.

In general $u_1 \neq 1 - u_2$. But it turns out that performing the γ_{12}-integration in expression eq.(116) produces a derivative of a Dirac δ-function $\delta(u_1 + u_2 - 1)$.

Expecting this, the two x-dependent factors in expression (119) are combined using identity (118), with integration variable v. Noting that the expression (121) is linear in u, and using identity (121) appropriately, the result for $\underline{\overline{X}}$ can be written as follows, using eq.(114)

$$\underline{\overline{X}} \;=\; \int\int du_1 du_2 \left(\frac{u_1}{1-u_1}\frac{u_2}{1-u_2}\right)^{-\frac{1}{2}\gamma_{12}} \\ Y(u_1, u_2 | x_4, \ldots, x_1)$$

with $\hat{\beta}_{12} = \hat{\delta}_{12} - \delta^0_{12}$ and

$$
\begin{aligned}
&Y(u_1,u_2|x_4,\ldots,x_1) = \qquad (123)\\
&\quad u_1^{\delta^0_{13}-\frac{1}{2}\hat{\beta}_{12}-1}(1-u_1)^{\delta^0_{14}-\frac{1}{2}\hat{\beta}_{12}-1}\\
&\quad u_2^{\delta^0_{24}-\frac{1}{2}\hat{\beta}_{12}-1}(1-u_2)^{\delta^0_{23}-\frac{1}{2}\hat{\beta}_{12}-1}\\
&\int dv v^{\delta^0_{13}+\delta^0_{14}-\hat{\beta}_{12}-1}(1-v)^{\delta^0_{23}+\delta^0_{24}-\hat{\beta}_{12}-1}\\
&\Gamma(\Delta_{12})\left(-\frac{1}{2}x_{05}(v,1-u_2)-\frac{1}{2}v(1-v)x_{12}^2-\right.\\
&\left.-\frac{1}{2}u_2(1-u_2)x_{34}^2-\frac{1}{2}v(u_1+u_2-1)(x_{13}^2-x_{14}^2)\right)^{-\Delta_{12}}
\end{aligned}
$$

with

$$
\begin{aligned}
\Delta_{12} &= \delta^0_{13}+\delta^0_{14}+\delta^0_{23}+\delta^0_{24}-2\hat{\beta}_{12}\\
&= d_1+d_2-2\hat{\delta}_{12}\\
&= d_3+d_4-2\hat{\delta}_{34}\ . \qquad (124)
\end{aligned}
$$

All integrations over u_1, u_2 and v run from 0 to 1. It is understood that $x_{05}(v,1-u_2) = x_0(v) - x_5(1-u_2)$ and $\hat{\beta}_{12} = \hat{\delta}_{12} - \delta^0_{12}$.

The factor $Y(u_1,u_2|x_4,\ldots,x_1)$ does not depend on γ_{12}.

The expression (116) for R involves

$$
\begin{aligned}
&(2\pi i)^{-1}\int_{-i\infty}^{i\infty} d\gamma_{12} P(\hat{\delta}_{12}|\gamma_{12})\underline{\overline{X}} = (2\pi i)^{-1}\int_{-i\infty}^{i\infty} d\gamma_{12}\\
&\int\int du_1 du_2 P(\hat{\delta}_{12}|\gamma_{12})\left(\frac{u_1}{1-u_1}\frac{u_2}{1-u_2}\right)^{-\frac{1}{2}\gamma_{12}}\\
&\quad Y(u_1,u_2|x_4,\ldots,x_1)
\end{aligned}
$$

The γ_{12}-integration can be performed. P being a polynomial in γ_{12}, we get a derivative of a Dirac δ-function $\delta(y_1+y_2)$ with $y_1 = -\frac{1}{2}\ln(u_1/(1-u_1))$ and $y_2 = -\frac{1}{2}\ln(u_2/(1-u_2))$..

The argument of the δ-function becomes 0 if and only if $u_2 = 1-u_1$. One may use the formula

$$\delta(f(x)) = |f'(x_0)|^{-1}\delta(x-x_0)\ ,$$

which is valid when $f(x)$ has a single zero at $x = x_0$. In our case the variable is u_1, and $f(u_1) = y_1 + y_2$, so that $f'(u_1)^{-1} = -2u_1(1-u_1)$. The result is

$$(2\pi i)^{-1} \int_{-i\infty}^{i\infty} \frac{1}{2} d\gamma_{12} P(\hat{\delta}_{12}|\gamma_{12}) \underline{\overline{X}} \tag{125}$$
$$= \int\int du_1 du_2 \; u_1(1-u_1) Y(u_1, u_2|x_4, \ldots, x_1)$$
$$P\left(\hat{\delta}_{12}|2u_1(1-u_1)\frac{\partial}{\partial u_1}\right)\delta(u_1+u_2-1)$$

This can be inserted into the expression (116) for R. Thanks to the prescription $u_2 = 1 - u_1$, which results from the δ-function, the x-dependent last factor in Y becomes a propagator. But before the prescription can be applied, the derivatives with respect to u_1 must be performed. This gives a sum of terms involving propagators whose exponents differs from $-\Delta_{12}$ by non-positive integers, and extra factors of $x_{13}^2 - x_{14}^2$, which tend to zero when $x_3 - x_4 \mapsto 0$.

Now we wish to obtain an asymptotic expansion as $x_{12} \equiv x_1 - x_2 \mapsto 0$ and $x_{13} \equiv x_3 - x_4 \mapsto 0$, keeping the two pairs of points x_1, x_2 and x_3, x_4 well separated. In this limit, x_{05} tends to a nonzero limit which is independent of v and u_2.

To obtain an asymptotic expansion for future use expand the x-dependent last factor in Y in powers of x_{12}^2 and x_{34}^2, using the expansion formula

$$\Gamma(\alpha)\,(A-\epsilon)^{-\alpha} = \sum_k \frac{1}{k!}\epsilon^k \Gamma(\alpha+k) A^{-\alpha-k} \tag{126}$$

We use this first for $\epsilon = \frac{1}{2}v(1-v)x_{12}^2$ and then again for $\epsilon = \frac{1}{2}u_2(1-u_2)x_{34}^2$.

The result is

$$
\begin{aligned}
&Y(u_1,u_2|x_4,\ldots,x_1) = \\
&\sum_{p=0}^{\infty}\sum_{q=0}^{\infty}\frac{(-)^{p+q}}{p!q!}\left(-\frac{1}{2}x_{12}^2\right)^p\left(-\frac{1}{2}x_{34}^2\right)^q \\
&\quad u_1^{\delta_{13}^0-\frac{1}{2}\hat{\beta}_{12}+q-1}(1-u_1)^{\delta_{14}^0-\frac{1}{2}\hat{\beta}_{12}+q-1} \\
&u_2^{\delta_{24}^0-\frac{1}{2}\hat{\beta}_{12}-1}(1-u_2)^{\delta_{23}^0-\frac{1}{2}\hat{\beta}_{12}-1} \\
&\quad\int dv\ v^{\delta_{13}^0+\delta_{14}^0-\hat{\beta}_{12}+p-1}(1-v)^{\delta_{23}^0+\delta_{24}^0-\hat{\beta}_{12}+p-1} \\
&\quad\Gamma(\Delta_{12}+p+q)\left(-\frac{1}{2}x_{05}(v,1-u_2)^2\right. \\
&\qquad\left.-\frac{1}{2}v(u_1+u_2-1)(x_{13}^2-x_{14}^2)\right)^{-\Delta_{12}-p-q}
\end{aligned}
\tag{127}
$$

$-R$ is obtained by multiplying expression (125) with the two propagators $\Gamma(\hat{\delta}_{12})\left(-\frac{1}{2}x_{12}^2\right)^{-\hat{\delta}_{12}}\Gamma(\hat{\delta}_{34})\left(-\frac{1}{2}x_{34}^2\right)^{-\hat{\delta}_{34}}$, inserting eq.(127) for Y, and performing the differentiations in P.

For future reference let us specialize to the case when the polynomial P is constant in γ_{12},

$$P(\hat{\delta}|\gamma_{12}) = r(\hat{\delta}) \in \mathbf{R}\ ,$$

and for equal dimensions $d_1 = d_2 = d_3 = d_4 = d$, whence

$$\hat{\delta}_{34} = \hat{\delta}_{12} = \hat{\delta} + d$$

by eq.(115).

The resulting asymptotic expansion for R is

$$
\begin{aligned}
&R(\hat{\delta}|x_4,\ldots,x_1) = -r(\hat{\delta})\sum_{p=0}^{\infty}\sum_{q=0}^{\infty}\frac{(-)^{p+q}}{p!q!} \\
&\Gamma(\hat{\delta}_{12})\left(-\frac{1}{2}x_{12}^2\right)^{-\hat{\delta}_{12}+p}\Gamma(\hat{\delta}_{12})\left(-\frac{1}{2}x_{34}^2\right)^{-\hat{\delta}_{12}+q} \\
&\quad\int du\int dv[u(1-u)]^{d-\hat{\delta}_{12}+q-1}[v(1-v)]^{d-\hat{\delta}_{12}+p-1} \\
&\quad\Gamma(2d-2\hat{\delta}_{12}+p+q)\left(-\frac{1}{2}x_{05}^2(v,u)\right)^{-2d+2\hat{\delta}_{12}-p-q}\ ,
\end{aligned}
\tag{128}
$$

where $x_0(v) = vx_1 + (1-v)x_2$, $x_5(u) = ux_3 + (1-u)x_4$ and $x_{05}(v,u) = x_0(v) - x_5(u)$.

11 Comparison of contributions to asymptotic expansions as $x_{12} \mapsto 0$ from OPE and from families of poles in the Mellin amplitude

We compare the contribution to the asymptotic expansion of the scalar 4-point Green function G_{0000} as $\mathbf{x}_{12} \mapsto 0$ and/or $\mathbf{x}_{34} \mapsto 0$ which come from a scalar field in the OPE with the contribution of the corresponding family of dynamical poles in the Mellin amplitude. The kinematical poles need not be considered since they just cancel disconnected parts, by their definition.

11.1 Summation of the contributions from a family of poles in the Mellin amplitude which corrresponds to a scalar field in the OPE

Consider the four-point function G_{0000} of four real fields ϕ^0 of dimension d.

As we have seen in section 8.1, a scalar field ϕ^k of dimension $\delta = h + c$ which contributes to the operator product expansion in the (12)-channel produces a family of poles in δ_{12} in the Mellin amplitude at positions $\delta^{(43)(21)} = \delta_0^n(\delta)$, whence

$$\hat{\delta}_{12}^n = d - \frac{1}{2}\delta - n, \qquad , n = 0, 1, ... \tag{129}$$

with residues as follows: For the scalar case, $\chi = [0, h+c]$, the polynomial $\mathfrak{P}_\chi^D$ is of degree 0 and equals $\prod_{n=1}^4 \Gamma(\delta_n)^{-1}$, with δ_i given by eqs.(94). Therefore the residue r_n of the n-th pole obtains from eq.(108)f as

$$r_n = \frac{const}{\Gamma(\delta_{12})\Gamma(\delta_{34})} \, \frac{(-)^n}{n!}\Gamma(-n-c) \; . \tag{130}$$

with

$$\begin{aligned} const &= -\frac{\sin \pi c}{\pi}|g_k^{00}|^2(\pi/2)^{-h} \\ &\quad \Gamma(\frac{1}{2}(h+c))^{-2}\Gamma(\frac{1}{2}(h+c))^{-2} \end{aligned} \tag{131}$$

In the expression for r_n, the arguments $\delta_{12} = \delta_{34}$ are determined by the position of the pole and the two inverse Γ-functions cancel against the first two Γ-functions in expression (128) for R.

The residues have the same sign for all n. [6]

The n-th pole makes a contribution R_n to the asymptotic expansion of the four-point function, which is given by eq.(128), with $\delta_0^n(\delta)$ substituted for $\hat{\delta}$, and

$$r(\hat{\delta}) = r_n.$$

We wish to sum these contribution over n.

To do so, we introduce new summation variables N,K in place of p, q by

$$\begin{aligned} n+p &= N \\ n+q &= K \end{aligned} \tag{132}$$

They run over $N \geq n, K \geq n$, but we may extend the summation from 0, because the extra contributions vanish, since $\frac{1}{p!} = \Gamma(N-n+1)^{-1} = 0$ when $N < n$, and similarly for K. Inserting expression (129), the sum of the contributions R_n comes out as

$$\begin{aligned} &\sum_n R(\delta_0^n(\delta)|x_4,...,x_1) = -const \\ &\sum_{N,K} S_{N,K}(c) \left(-\frac{1}{2}x_{12}^2\right)^{-d+\frac{1}{2}\delta+K} \left(-\frac{1}{2}x_{34}^2\right)^{-d+\frac{1}{2}\delta+N} \\ &\quad \int\int dudv[u(1-u)]^{\frac{1}{2}\delta+K-1}[v(1-v)]^{\frac{1}{2}\delta+N-1} \\ &\Gamma(\delta+N+K)\left(-\frac{1}{2}x_{05}^2(v,u)\right)^{-\delta-N-K} \end{aligned} \tag{133}$$

where $\delta = h + c$ is the dimension of the field ϕ^k in the OPE and

$$S_{N,K} = \sum_n \frac{(-)^n}{n!}\Gamma(-c-n)\frac{(-)^{N+K}}{(N-n)!(K-n)!} \tag{134}$$

[6] This implies that the contribution from individual poles is not positive. Canceling a pole with $n > 0$ would require adding a ghost field of dimension $\delta + 2n$ which contributes to the OPE

$S_{N,K}$ is manifestly symmetrical in N and K. We may therefore assume without generality that $N \geq K$. The sum can be evaluated by writing

$$\frac{\Gamma(-c-n)}{\Gamma(N-n+1)} = \frac{B(-c-n, c+N+1)}{\Gamma(c+N+1)} \ ,$$

inserting the standard integral representation for the Beta-function and interchanging sum and integral.The result can be written in the form

$$S_{N,K} = \frac{-\pi}{\sin \pi c} \frac{1}{K!N!} \frac{(-)^K}{\Gamma(c+N+1)} \frac{\Gamma(-c-K)}{\Gamma(-c-K-N)} \tag{135}$$

for $N \geq K$.

11.2 Short distance behavior from the OPE

Consider the Wightman 4-point function of four scalar fields $\phi^{i_4}, ..., \phi^{i_1}$ of dimensions $d_{i_4}, ..., d_{i_1}$.

According to eq.(19), the operator product expansion takes the form

$$\begin{aligned} W_{\bar{i_4},\bar{i_3},i_2,i_1}(x_4, ..., x_1) &= \sum_k c_k \bar{g}_k^{i_3 i_4} g_k^{i_2 i_1} \\ &\quad \mathfrak{W}_{\chi_k,\bar{i_4},\bar{i_3},i_2,i_1}(x_4, ..., x_1) \ ; \end{aligned} \tag{136}$$

where $\bar{g}$ is the complex conjugate of g.

The term labeled by k represents the contribution of field ϕ^k with Lorentz spin and dimension given by $\chi_k = [l_k, d_k]$; $c_k > 0$ fixes the normalization of its two point functions. The c_k and coupling constants $g_{\cdot\cdot}^{\cdot}$ are the dynamical quantities which need to be determined in order to construct a theory, and $\mathfrak{W}$ are kinematically determined quantities.

There is freedom of choosing normalization factors.

According to eq.(25),

$$\begin{aligned} \mathfrak{W}_{l,m,j,i}(x_4, ..., x_1) &= \mathfrak{Q}(\chi_k, p; \chi_l, x_4, \chi_m, x_3) \\ &\quad \mathfrak{V}(\chi_k, z, \chi_j, x_2, \chi_i, x_1)|_{z=0} \ . \end{aligned} \tag{137}$$

with $p = -i\nabla_z$. It is understood that $c_-^{43} = d_l - d_m$ etc.

The kinematically determined quantity $\mathfrak{W}$ given by eq. (137) is boundary value of an analytic function, whose domain of analyticity (the tube) includes Euklidean points. Upon substitution of $\mathfrak{Q}^u$ and $\mathfrak{V}^u$ for $\mathfrak{Q}$ and $\mathfrak{V}$, I denote this

quantity by $(2\pi)^{-D} I_l^u$, where l is the Lorentz spin of the field ϕ^k in the OPE. I will evaluate this quantity for $l = 0$, thereby computing the contribution of a scalar field in the OPE to the short distance behavior.

Inserting the formulas (77) for $\mathfrak{V}^u$, and (175) for the partial Fourier transform of $\mathfrak{Q}^u$ into eq.(137), the factor $e^{ip[ux_4+(1-u)x_3]}$ acts as a translation operator which shifts z form 0 to $x_5 = ux_4 + (1-u)x_3$, and p^2 acts as $-\Box_5$. Inserting the power series expansion of J_c, one obtains

$$\begin{aligned} I_0^u &= \frac{-1}{\Gamma(\frac{1}{2}(h-c)+c_-^{43})\Gamma(\frac{1}{2}(h-c)-c_-^{43})} \\ &\quad \sum_{N=0}^{\infty} \frac{(-1)^N}{N!\Gamma(c+N+1)} 2^{-N} \\ &\quad (-\frac{1}{2}x_{43}^2)^{-\frac{1}{2}(h-c)-c_+^{43}+N} (-\frac{1}{2}x_{12}^2)^{-\frac{1}{2}(h-c)-c_+^{12}} (-\Box_5)^N \\ &\quad \int_0^1 du\ u^{\frac{1}{2}(h+c)+c_-^{43}+N-1}\ (1-u)^{\frac{1}{2}(h+c)-c_-^{43}+N-1} \\ &\quad (-\frac{1}{2}x_{25}^2)^{-\frac{1}{2}(h+c)+c_-^{12}} (-\frac{1}{2}x_{15}^2)^{-\frac{1}{2}(h+c)-c_-^{12}} \\ x_5 &\equiv ux_4 + (1-u)x_3 \end{aligned}$$

Now the two propagators in the last line are combined into a single propagator by use of the standard identity (118), introducing an auxiliary variable of integration v. Defining

$$x_0(v) = vx_2 + (1-v)x_1, \tag{138}$$

and $x_{05} = x_0 - x_5$, the new propagator can be rewritten with the help of the identity

$$vx_{20}^2 + (1-v)x_{10}^2 = x_{05}^2 + v(1-v)x_{12}^2\ .$$

and comes out proportional to

$$\begin{aligned} &\Gamma(h+c)(-\frac{1}{2}x_{05}^2 - \frac{1}{2}v(1-v)x_{12}^2)^{-h-c} = \\ &\sum_{K=0}^{\infty} \frac{1}{K!}(\frac{1}{2}v(1-v)x_{12})^2)^K \Gamma(h+c+K)(-\frac{1}{2}x_{05}^2)^{-h-c-K} \end{aligned} \tag{139}$$

Finally, the differentiation $\Box_5^N$ is carried out with the help of the formula

$$(\Box_5)^N\Gamma(\delta)\left(-\frac{1}{2}x_{05}^2\right)^{-\delta} = \tag{140}$$
$$2^N\frac{\Gamma(-\delta+h)}{\Gamma(-\delta+h-N)}\Gamma(\delta+N)\left(-\frac{1}{2}x_{05}^2\right)^{-\delta-N}$$

Collecting everything and noting that $\delta_\chi = \frac{1}{2}(h+c)$ produces the final result

$$\begin{aligned}
I_0^u &= \frac{-1}{\Gamma(\frac{1}{2}(h-c)+c_-^{43})\Gamma(\frac{1}{2}(h-c)-c_-^{43})} \\
&\quad \frac{1}{\Gamma(\frac{1}{2}(h+c)+c_-^{12})\Gamma(\frac{1}{2}(h+c)-c_-^{12})} \\
&\quad \sum_{N=0}^{\infty}\sum_{K=0}^{\infty}\frac{(-1)^K}{K!N!}\frac{\Gamma(-c-K)}{\Gamma(c+N+1)\Gamma(-c-K-N)} \\
&\quad \left(-\frac{1}{2}x_{43}^2\right)^{-\frac{1}{2}(h-c)-c_+^{43}+N}\left(-\frac{1}{2}x_{12}^2\right)^{-\frac{1}{2}(h-c)-c_+^{12}+K} \\
&\quad \int_0^1 du\ u^{\frac{1}{2}(h+c)+c_-^{43}+N-1}(1-u)^{\frac{1}{2}(h+c)-c_-^{43}+N-1} \\
&\quad \int_0^1 dv\ v^{\frac{1}{2}(h+c)-c_-^{12}+K-1}(1-v)^{\frac{1}{2}(h+c)+c_-^{12}+K-1} \\
&\quad \Gamma(h+c+K+N)\left(-\frac{1}{2}x_{05}^2\right)^{-h-c-N-K} \qquad . \qquad (141) \\
x_0 &= ux_1+(1-u)x_2 \\
x_5 &= vx_3-(1-v)x_4
\end{aligned}$$

and $x_{05} = x_0 - x_5$. According to eqs.(19,23), the contribution of a real scalar field ϕ^k of dimension $d_k = h + c$ in the OPE to the 4-point function $W_{i_4,\ldots,i_1}$ equals

$$(2\pi)^{-D}\bar{g}_k^{i_3 i_4}g_k^{i_2 i_1}\mathbf{c}_k I_0^u, \tag{142}$$

where $\mathbf{c}_k$ is determined by the normalization of the two-point function of ϕ^k.

11.3 Comparison with the expansion from the Mellin representation

We now compare the contribution (142) of a scalar field to the OPE (85) of the Euklidean 4-point function with the sum (133) of the contribution of the corresponding family of poles in the Mellin representation, with expressions (135) for $S_{N,K}$ and (131) for $const$, for $c_-^{12} = c_-^{43} = 0$.

We see that both expressions agree, up to a constant factor which may be attributed to the normalization of two point functions.

The formulas obtained by summation of contributions from a family of Mellin poles generalize to arbitrary dimensions of the scalar fields and agreement is again obtained.

Acknowledgment

I would like to thank Volker Schomerus and Yuri Pis'mak for stimulating discussions and for reading preliminary versions of the manuscript. Ivan Todorov kindly informed me that Bakalov and Nikolov had independently studied dimensional induction [3].

12 Appendix. Symanziks n-star integration formula in Euklidean space

In ref.[70], Symanzik presented a formula which gives a Mellin representation of the integral of $n \geq 3$ conformal invariant scalar propagators over D-dimensional Euklidean space under conditions which guarantee the conformal invariance of the result.

In the following, all variables $\mathbf{x}_i$ will be in Euklidean space, and

$$\mathbf{x}_{ij} = \mathbf{x}_i - \mathbf{x}_j \ .$$

The formula reads as follows

$$(\pi)^{-\frac{1}{2}D}\int d^D\mathbf{x}_0\prod_{i=1}^{n}\left(\mathbf{x}_{0i}^2\right)^{-\delta_i}\Gamma(\delta_i)= \tag{143}$$
$$(2\pi i)^{-\frac{1}{2}n(n-3)}\int_{-i\infty}^{i\infty}...\int_{-i\infty}^{i\infty}ds_1...ds_{\frac{1}{2}n(n-3)}$$
$$\prod_{1\le i<j\le n}\Gamma(\delta_{ij}(s))(\mathbf{x}_{ij}^2)^{-\delta_{ij}(s)}$$

provided

$$0<\Re e\delta_i<\frac{1}{2}D \tag{144}$$

for all i, and ,

$$\sum_i\delta_i=D. \tag{145}$$

Integration is over the hyper-surface of all solutions (with fixed real part) $\delta_{ij}=\delta_{ji}$, $(i,j=1...n, i\neq j)$ of the system of equations

$$\sum_{j\neq i}\delta_{ij}=\delta_i, \tag{146}$$

parametrized by $\frac{1}{2}n(n-3)$ imaginary variables $s_1,...,s_{\frac{1}{2}n(n-3)}$ as follows.

Choose a particular solution δ_{ij}^0 of eqs.(146) subject to the inequalities (144), and real coefficients $c_{ij,k}=c_{ji,k}$ which obey

$$c_{ii,k}=0,\quad \sum_{j\neq i}c_{ij,k}=0. \tag{147}$$

Then

$$\delta_{ij}=\delta_{ij}^0+\sum_{k=1}^{\frac{1}{2}n(n-3)}c_{ij,k}s_k. \tag{148}$$

To obtain the proper normalization of the measure on the hyper-surface it is demanded that the $(\frac{1}{2}n(n-3))^2$ coefficients $c_{ij,k}$ with $2\le i<j\le n$, excepting $c_{23,k}$, which may be taken as the independent ones, must satisfy

$$|det\ c_{ij,k}|=1. \tag{149}$$

If $n = 3$ there is no integration over s to be performed, and eq.(143) is a star-triangle relation.

Our main interest will be in the case $n = 4$ and its generalization with spin. The general solution of eqs.(146) takes the form

$$\delta_{ij} = \delta^0_{ij} + \beta_{ij}, \tag{150}$$

with $\beta_{ij} = \beta_{ji}$ and

$$\beta_{34} = \beta_{12},\ \beta_{24} = \beta_{13},\ \beta_{14} = \beta_{23} \tag{151}$$

$$\beta_{12} + \beta_{13} + \beta_{23} = 0 \tag{152}$$

Choosing $s_1 = \beta_{12}, s_2 = \beta_{13}$ the above conditions on the coefficients $c_{ij,k}$ are fulfilled, and similarly for any other choice of two out of the three variables $\beta_{12}, \beta_{13}, \beta_{23}$. Thus we may write

$$ds_1 ds_2 = d^2\beta = d\beta_{12} d\beta_{13} = ... \tag{153}$$

12.1 Generalization of the 4-star formula

In the computation of the Mellin representation $\mathfrak{M}_\chi$ of a Euklidean partial wave, a generalization of the Symanzik 4-star formula is needed which includes a factor $Y_l^D(\cos\theta)$ where

$$\cos\theta = \frac{\lambda\mu}{|\lambda||\mu|} \tag{154}$$

is the cosine of the angle between the Euklidean D-vectors

$$\lambda = 2\left(\frac{\mathbf{x}_{10}}{\mathbf{x}^2_{10}} - \frac{\mathbf{x}_{20}}{\mathbf{x}^2_{20}}\right) \tag{155}$$

and

$$\mu = 2\left(\frac{\mathbf{x}_{40}}{\mathbf{x}^2_{40}} - \frac{\mathbf{x}_{30}}{\mathbf{x}^2_{30}}\right)\ . \tag{156}$$

In applications, $Y_l^D(\cos\theta)$ is the zonal spherical function in D dimensions

Its presence reflects the fact that the quantum numbers of a field of Lorentz spin l are exchanged between legs $1, 2$ and $3, 4$. But the result holds for arbitrary polynomials of the form

$$Y_l^D(t) = \sum_{k=0}^{E[l/2]} a_k^l(D) t^{l-2k}\ , \tag{157}$$

i.e. which involves even powers of t for even l, and odd powers for odd l. $E[l/2]$ is the largest integer $\leq l/2$,.

Given the external dimensions δ_i, the result of the Symanzik integration formula involves an integration over two independent variables s_1, s_2 - e.g. δ_{12} and δ_{13} - which determine the six $\delta_{ij} = \delta_{ji}$ subject to the constraint

$$\sum_j \delta_{ij} = \delta_i \ . \tag{158}$$

We may equally well consider functions of δ_i and these variables as functions of the six δ_{ij}. Because validity of the Symanzik integration formula presupposes $\sum \delta_i = D$, they obey $\sum_i \sum_j \delta_{ij} = D$, but the definition of $\mathfrak{P}_l^D$ will make sense for arbitrary δ_{ij}.

The result will involve functions $\mathfrak{P}_l^D(\{\delta_{ij}\})$ which are polynomials of degree l in the independent variables among the δ_{ij}, for given δ_i, and such that $\mathfrak{P}_l^D(\{\delta_{ij}\}) \prod_{i=1}^4 \Gamma(\delta_i + \frac{l}{2})$ is a polynomial also in the δ_i. The only explicit dependence on D, for given $\{\delta_{ij}\}$, is through the expansion coefficients $a_k^l(D)$ of $Y_l^D(t)$. Explicitly, the polynomials are defined as follows

$$\mathfrak{P}_l^D(\{\delta_{ij}\}) = \sum_{k=0}^{E[l/2]} a_k^l(D) \ (\delta_{12} - \frac{1}{2}l)_k (\delta_{34} - \frac{1}{2}l)_k \mathfrak{P}_{l-2k}(\{\delta_{ij}\}) \tag{159}$$

where $\mathfrak{P}_m$ depends neither explicitly on D nor on l and is explicitly defined as follows.

The definition involves a sum over nonnegative integers $k_{ij} = k_{ji}$ attached to the four links $(ij) = (13), (14), (23), (24)$. A sum or product over these four links will be denoted by $\sum'$ and $\prod'$ respectively. In the definition of $\mathfrak{P}_m$, the variables are subject to the constraint $\sum' k_{ij} = m$.

The definition of $\mathfrak{P}_m$ is as follows.

$$\mathfrak{P}_m(\{\delta_{ij}\}) = 2^{-m} m! \sum_{\{k_{ij}\}: \sum' k_{ij} = m} \frac{(-)^{k_{14}+k_{23}}}{\prod' k_{ij}!} \prod{}' (\delta_{ij})_{k_{ij}} \prod_{n=1}^4 \Gamma\left(\delta_n - \frac{m}{2} + \sum_j k_{jn}\right)^{-1} \ . \tag{160}$$

Here as everywhere we use the notation

$$(\alpha)_k = \Gamma(\alpha + k)/\Gamma(\alpha) = \alpha(\alpha+1)...(\alpha+k-1) \ ,$$

and δ_n is to be read as $\sum_j \delta_{nj}$.

For fixed δ_i, $\mathfrak{P}_m$ is a polynomial in the remaining variables, and $\prod_{n=1}^4 \Gamma(\delta_n + \frac{l}{2})\mathfrak{P}_{l-2k}(\{\delta_{ij}\})$ is a polynomial also in the δ_n.

The generalization of the Symanzik 4-star formula reads as follows. Given δ_i, $i = 1...4$ subject to the constraint $\sum_i \delta_i = D$, let $\cos\theta$ be defined through eqs.(154)ff. Then

$$\begin{aligned} S_l &\equiv \pi^{-\frac{1}{2}D} \int d^D\mathbf{x}_0 Y_l^D(\cos\theta) \prod_{i=1}^4 (\mathbf{x}_{i0}^2)^{-\delta_i} && (161) \\ &= (2\pi i)^{-2} \int\int ds_1 ds_2 \mathfrak{P}_l^D(\{\delta_{ij}\}) && (162) \\ &\quad \Gamma\left(\delta_{12} - \frac{1}{2}l\right)(\mathbf{x}_{12}^2)^{-\delta_{12}} \Gamma\left(\delta_{34} - \frac{1}{2}l\right)(\mathbf{x}_{34}^2)^{-\delta_{34}} \\ &\quad \prod{}' \Gamma(\delta_{ij})(\mathbf{x}_{ij}^2)^{-\delta_{ij}} \ . \end{aligned}$$

The integrations are as in the ordinary Symanzik 4-star formula. It corresponds to the special case $l = 0$, with

$$\mathfrak{P}_0^D(\{\delta_{ij}\}) = \prod_{i=1}^4 \Gamma(\delta_i)^{-1}.$$

12.2 Derivation of the generalized 4-star formula

One computes

$$\begin{aligned} \lambda^2 &= 4\frac{\mathbf{x}_{12}^2}{\mathbf{x}_{10}^2\mathbf{x}_{20}^2} && (163) \\ \mu^2 &= 4\frac{\mathbf{x}_{34}^2}{\mathbf{x}_{30}^2\mathbf{x}_{40}^2} && (164) \\ \lambda\mu &= 2\left(\frac{\mathbf{x}_{13}^2}{\mathbf{x}_{10}^2\mathbf{x}_{30}^2} + \frac{\mathbf{x}_{24}^2}{\mathbf{x}_{20}^2\mathbf{x}_{40}^2} - \frac{\mathbf{x}_{14}^2}{\mathbf{x}_{10}^2\mathbf{x}_{40}^2} - \frac{\mathbf{x}_{23}^2}{\mathbf{x}_{20}^2\mathbf{x}_{30}^2}\right) \ . && (165) \end{aligned}$$

The multinomial theorem asserts

$$\begin{aligned} (\lambda\mu)^{l-2k} &= 2^{l-2k} \sum_{\{k_{ij}\}:\sum' k_{ij} = l-2k} \frac{(l-2k)!\ (-)^{k_{14}+k_{23}}}{\prod' k_{ij}!} \\ &\quad \prod{}' (\mathbf{x}_{ij}^2)^{k_{ij}} (\mathbf{x}_{i0}^2\mathbf{x}_{j0}^2)^{-k_{ij}} \end{aligned}$$

Inserting the power series expansion (157) of Y_l^D, one obtains

$$\begin{aligned}
&\pi^{\frac{1}{2}D} S_l = \\
&\int d^D\mathbf{x}_0 \sum_{k=0}^{E[l/2]} a_k^l(D)(\lambda^2\mu^2)^{-\frac{1}{2}l+k}(\lambda\mu)^{l-2k} \prod (\mathbf{x}_{i0})^{-\delta_i} \\
= &\int d^D\mathbf{x}_0 \sum_{k=0}^{E[l/2]} a_k^l(D) 2^{-l+2k}(l-2k)!(\mathbf{x}_{12}^2\mathbf{x}_{34}^2)^{-\frac{1}{2}l+k} \\
&\sum_{\{k_{ij}\}:\sum' k_{ij}=l-2k} \frac{(-)^{k_{14}+k_{23}}}{\prod' k_{ij}!} {\prod}' (\mathbf{x}_{ij}^2)^{k_{ij}} \prod_i (\mathbf{x}_{0i}^2)^{-\delta_i'} \ , \qquad (166)
\end{aligned}$$

with

$$\delta_i' = \delta_i + \sum_j k_{ij} - \frac{1}{2}l + k \ . \qquad (167)$$

One verifies that $\sum \delta_i' = D$. Therefore the integral can be performed with the help of the Symanzik 4-star formula. It will involve exponents δ_{ij}' subject to the constraint $\sum_j \delta_{ij}' = \delta_i'$, and produce factors

$$\prod_{n=1}^4 \Gamma(\delta_n')^{-1} \prod_{ij} \Gamma(\delta_{ij}')(\mathbf{x}_{ij}^2)^{-\delta_{ij}'}$$

which multiply the preexisting factors $(\mathbf{x}_{12}^2\mathbf{x}_{34}^2)^{-\frac{1}{2}l+k} \prod'(\mathbf{x}_{ij}^2)^{k_{ij}}$. Transform to new variables

$$\begin{aligned}
\delta_{ij} &= \delta_{ij}' - k_{ij} \ , \quad (ij) \neq (12), (34), \\
\delta_{12} &= \delta_{12}' + \frac{1}{2}l - k \ , \\
\delta_{34} &= \delta_{34}' + \frac{1}{2}l - k \ . \qquad (168)
\end{aligned}$$

As a consequence of the constraint on the δ_{ij}', they satisfy the constraint $\sum_j \delta_{ij} = \delta_i$.

It follows that

$$\begin{aligned}
\Gamma(\delta_{ij}') &= \Gamma(\delta_{ij})(\delta_{ij})_{k_{ij}} \ , \quad (ij) \neq (12), (34) \\
\Gamma(\delta_{12}') &= \Gamma(\delta_{12} - \frac{1}{2}l)(\delta_{12} - \frac{1}{2}l)_k \\
\Gamma(\delta_{34}') &= \Gamma(\delta_{34} - \frac{1}{2}l)(\delta_{34} - \frac{1}{2}l))_k
\end{aligned}$$

Inserting this and carrying out the sum over the k_{ij} with the help of the definition of $\mathfrak{P}_m$, and the sum over k with the help of the definition of $\mathfrak{P}_l^D$ one obtains the final result, eq.(162).

The polynomial properties of $\mathfrak{P}_m$ and therefore of $\mathfrak{P}_l^D$ for fixed δ_i are obvious from the fact that $(\alpha)_k = \Gamma(\alpha+k)/\Gamma(\alpha)$ is a polynomial in α of degree k.

$\prod_{n=1}^4 \Gamma(\delta_n + \frac{l}{2})\mathfrak{P}_{l-2k}(\{\delta_{ij}\})$ is a polynomial also in δ_n because $\Gamma(\delta_n + \frac{l}{2})\Gamma(\delta_n - \frac{l}{2} + k + \sum_j k_{jn})^{-1}$ is a polynomial. This is so because $-l+k+\sum_j k_{jn}$ is an integer $\leq -k$ since $\sum_j k_{jn} \leq l - 2k$.

13 Appendix: 3-point vertex functions and coefficients in the operator product expansion

13.1 Notations

In $D = 2h$ space time dimensions, let $\chi = [l, \delta = h + c]$. Introduce

$$\tilde{\chi} = [\bar{l}, h - c], \tag{169}$$

$$\delta_\chi = \frac{1}{2}(h + c - l) \tag{170}$$

$$\delta_{\tilde{\chi}} = \frac{1}{2}(h - c - l) \tag{171}$$

$\bar{l}$ is the conjugate representation to l which is the same as l for symmetric tensor representations.

I also use the shorthand

$$(\alpha)_k = \frac{\Gamma(\alpha+k)}{\Gamma(\alpha)} = \alpha(\alpha+1)\dots(\alpha+k-1)\ .$$

13.2 The differential operators D_l

Following ref.[14], a convenient alternative expression for the conformal invariant 3-point function $\mathfrak{V}$ can be written down which uses a l-th order differential operator D_l. It acts on functions of x_1 and x_2. It is homogeneous in the complex light-like D-vector $\mathfrak{z}$ of degree l.

$D_l(\delta_\chi + c_-, \mathfrak{z}\nabla_1, \delta_\chi - c_-, \mathfrak{z}\nabla_2)$ is defined by

$$\begin{aligned} & D_l(a,\alpha;b,\beta) = \\ & \sum_{k=0}^{l} \left(\begin{array}{c} l \\ k \end{array} \right) (a+k)_{l-k}(b+l-k)_k(-\alpha)^k\beta^{l-k} \\ = \quad & (a)_l(\alpha+\beta)^l F\left(a+b+l-1,-l;a;\frac{\alpha}{\alpha+\beta}\right) \end{aligned} \tag{172}$$

where $F = {}_2F_1$ is the hypergeometric function.

The following identity is valid, and easily checked by use of the binomial theorem

$$\begin{aligned} & (a)_l(b)_l \left(\frac{2}{x_{13}^2}\right)^a \left(\frac{2}{x_{23}^2}\right)^b (\lambda\cdot\mathfrak{z})^l \\ = \quad & D_l(a,\mathfrak{z}\cdot\nabla_1;b,\mathfrak{z}\cdot\nabla_2)\left(\frac{2}{x_{13}^2}\right)^a \left(\frac{2}{x_{23}^2}\right)^b \end{aligned} \tag{173}$$

with $\lambda = 2\left(\frac{x_{13}}{x_{13}^2} - \frac{x_{23}}{x_{23}^2}\right)$

13.3 The un-amputated 3-point vertex function with one spinning leg

Let $\chi_1 = [0, d_1 = h + c_1]$, $\chi_2 = [0, d_2 = h + c_2]$, $\chi = [l, h + c]$, and $c_+ = \frac{1}{2}(c_1 + c_2)$, $c_- = \frac{1}{2}(c_1 - c_2)$, and retain the definition $\delta_\chi = \frac{1}{2}(h + c - l)$ of subsection 13.1 Using the identity (173), expression (77) for the three-point function can be written in the alternative form

$$\begin{aligned} & \mathfrak{V}^u(\mathbf{x}_0,\chi,\mathbf{x}_1,\chi_1,\mathbf{x}_2,\chi_2) = \\ & (2\pi)^{-h}\frac{1}{(\delta_\chi+c_-)_l(\delta_\chi-c_-)_l}\left(-\frac{2}{x_{12}^2}\right)^{\frac{1}{2}(h-c+l)+c_+} \\ & D_l(\delta_\chi+c_-,\mathfrak{z}\nabla_1,\delta_\chi-c_-,\mathfrak{z}\nabla_2) \\ & \left[\left(\frac{-2}{x_{10}^2}\right)^{\frac{1}{2}(h+c-l)+c_-}\left(\frac{-2}{x_{20}^2}\right)^{\frac{1}{2}(h+c-l)-c_-}\right] \end{aligned} \tag{174}$$

This involves a Minkowski space scalar product so that $-x_{12}^2 > 0$ when x_{12} is space-like or Euklidean. The 3-point functions in Euklidean space and

in Minkowski space are obtained from each other by analytic continuation. $i\epsilon$-prescriptions as appropriate for Wightman functions are understood in Minkowski space. They follow from the spectrum condition.

13.4 The coefficient in the operator product expansion

From the formula (2.34) of [14] for the *scalar-leg-amputated* coefficient Q_- the un-amputeted coefficient is obtained by changing $c_1, c_2 \mapsto -c_1, -c_2$, assuming normalization conditions so that the scalar two-point function in momentum space for a field of dimension $h+c$ is $\left(\frac{1}{2}p^2\right)^c$, and rewriting the normalization factor in terms of $N_l(c_+, c_-, -c)$ by use of eq.(3.42) of ref. [11] . Using the notation of subsection 13.1, the result is as follows, for the partial Fourier transform

$$\begin{aligned}
&\mathfrak{Q}^u(p, \tilde{\chi}; x_1, \chi_1, x_2, \chi_2) = \qquad (175)\\
&-(2\pi)^{-h} \frac{1}{\Gamma(\frac{1}{2}(h-c+l)+c_-)\Gamma(\frac{1}{2}(h-c+l)-c_-)}\\
&\left(-\frac{1}{2}x_{12}^2\right)^{-\frac{1}{2}(h+c+l)-c_+} D_l(\delta_{\tilde{\chi}} + c_-, \mathfrak{z}\nabla_1, \delta_{\tilde{\chi}} - c_-, \mathfrak{z}\nabla_2)\\
&\int du [u(1-u)]^{\frac{h}{2}-1} \left(\frac{1-u}{u}\right)^{-c_-} \left(\frac{-x_{12}^2}{p^2}\right)^{\frac{l}{2}+\frac{c}{2}}\\
&J_{l+c}\left([-x_{12}^2 p^2 u(1-u)]^{\frac{1}{2}}\right)\ e^{ip[ux_1+(1-u)x_2]}
\end{aligned}$$

This is an entire function of p. In Euklidean space, the sign of the scalar product is reversed, so that $\mathbf{p}^2 = -\mathbf{p}^2 < 0$ and $x_{12}^2 = -\mathbf{x}_{12}^2 < 0$. One can use that $z^{-\nu} I_\nu(z\rho) = (e^{i\pi/2}z)^{-\nu} J_\nu(e^{i\pi/2}z\rho)$ for real ρ in order to express the result in Euklidean space in terms of the modified Bessel function I_{l+c}.

References

[1] Ofer Aharony, Steven S. Gubser, Juan Martin Maldacena, Hirosi Ooguri, and Yaron Oz. Large N field theories, string theory and gravity. *Phys. Rept.*, 323:183–386, 2000.

[2] D.P. Arovas, J.R. Schrieffer, F. Wilczek, and A. Zee. Statistical mechanics of anyons. *Nucl. Phys. B*, 251:117–126, 1985.

[3] B. Bakalov and N Nikolov. *(in preparation)*, 2009.

[4] B. Bakalov, N.M. Nikolov, K.H. Rehren, and I. Todorov. Infinite dimensional Lie algebra in 4d conformal quantum field theory. *J. Phys. A*, 41:194002, 2008.

[5] V. Bargmann and I.T. Todorov. Spaces of analytic functions on a complex cone as carriers for the symmetric tensor representations of $SO(n)$. *J. Math. Phys.*, pages 1141–1148, 1977. reviewed in [13], Appendix A.5.

[6] B. Basso, G.P. Korchemsky, and J. Kotański. Cusp anomalous dimension in maximally supersymmetric Yang Mills theory. *Phys. Rev. Letters*, 100:091601, 2008. eprint hep-th/07083933.

[7] Niklas Beisert. The dilatation operator of N = 4 super Yang-Mills theory and integrability. *Phys. Rept.*, 405:1–202, 2005.

[8] Curtis G. Callan and David J. Gross. Bjorken scaling in quantum field theory. *Phys. Rev.*, D8:4383–4394, 1973.

[9] Eric D'Hoker and Daniel Z. Freedman. Supersymmetric gauge theories and the AdS/CFT correspondence. In *TASI-lectures Boulder 2001: Strings,Branes and EXTRA dimensions*, pages 3–158, 2002.

[10] P.A.M. Dirac. *Ann. Math.*, 37:429, 1936.

[11] V.K. Dobrev, G. Mack, V.B. Petkova, S.G. Petrova, and I.T. Todorov. On Clebsch Gordan expansions for the Lorentz group in n dimensions. *Reports on Mathematical Physics*, 9:219, 1976.

[12] V.K. Dobrev, G. Mack, V.B. Petkova, S.G. Petrova, and I.T. Todorov. *Harmonic analysis on the n-dimensional Lorentz group and its applications to conformal quantum field theory*, volume 63 of *Lecture notes in physics*. Springer Verlag, 1977.

[13] V.K. Dobrev, G. Mack, V.B. Petkova, S.G. Petrova, and I.T. Todorov. *Harmonic analysis on the n-dimensional Lorentz group and its applications to conformal quantum field theory*, volume 63 of *Lecture notes in physics*. Springer Verlag, 1977.

[14] V.K. Dobrev, V.B. Petkova, S.G. Petrova, and I.T. Todorov. Dynamical derivation of operator-product expansions in Euclidean conformal quantum field theory. *Phys. Rev. D*, 13:887–912, 1976.

[15] J. Erdmenger and H. Osborn. Conserved currents and the energy momentum tensor in conformally invariant theories for general dimensions. *Nucl. Phys. B*, 483:431–474, 1997.

[16] S. Ferrara, R. Gatto, A.F. Grillo, and G. Parisi. The shadow operator formalism for conformal algebra. vacuum expectation values and operator products. *Lettere Nuovo Cimento*, 4:115, 1972.

[17] S. Ferrara, A.F. Grillo, and R. Gatto. *Conformal algebra in space-time and operator product expansions*, volume 67 of *Springer Tracts in Modern Physics*. Springer Verlag, 1973.

[18] S. Ferrara, A.F. Grillo, and R. Gatto. Tensor representations of conformal algebra amd conformally covariant operator product expansions. *Ann.Phys.(N.Y.)*, 76:161, 1973.

[19] S. Ferrara, A.F. Grillo, G. Parisi, and R. Gatto. Covariant expansion of the conformal four-point function. *Nucl. Phys. B*, 49:77, 1972.

[20] E.S. Fradkin. The quantum theory of fields i. *Zh. Exp. Teor. Fis.*, 29:121, 1955. in Russian. Engl. Translation: JETP 2 (1956) 148.

[21] E.S. Fradkin. Green functions in a quantum field theory and quantum statistics. In *Quantum field theory and hydrodynamics*, volume 29 of *Trudy FIAN*, Moscow, 1965. in Russian. Engl. translation: Plenum Press, New York 1967.

[22] E.S. Fradkin and M.Ya. Palchik. Method of solving conformal models in d-dimensional space. secondary fields in $d > 2$ and the solution of two-dimensional models. *Int. J. Mod. Phys. A*, 13:4837–4888, 1998.

[23] P.A. Frampton. *Dual Resonance Models and Superstrings*. World Scientific, 1986.

[24] K. Fredenhagen, B. Schroer, and K.H. Rehren. Superselection sectors with braid group statistics and exchange algebras, i: General theory. *Commun. Math. Phys.*, 125:201–226, 1989.

[25] P.G.O. Freund. Superstrings from 26 dimensions? *Physics Letters*, 151B:387–390, 1985.

[26] S. Fubini and G. Veneziano. Duality in operator formalism. *Nuovo Cimento*, 67A:29, 1970.

[27] M. Gel'fand and S. G. Gindikin. Complex manifolds whose skeletons are semisimple real Lie groups, and analytic discrete series of representations. *Funct. Anal. Appl.*, 7:258–265, 1978. Translation of *Funkts. Anal. Prilozh.*, 11(4):19-27, 1977.

[28] H. Georgi. Unparticle physics. *Phys. Rev. Letters.*, 98:221601, 2007.

[29] J. Glimm and A. Jaffe. *Quantum Physics. A Functional Integral Point of View*. Springer Verlag, 1981.

[30] I.S. Gradshteyn and I.M. Ryzhik. *Table of integrals, series, and products*. Academic Press, 1965.

[31] M.B. Green and J.H. Schwarz. Anomaly cancellation in supersymmetric $d = 10$ gauge theory and superstring theory. *Phys. Letters*, 149B:117–122, 1984.

[32] D.J. Gross, J.A. Harvey, E. Martinec, and R. Rohm. Heterotic string. *Phys. Rev. Letters*, 54:502–505, 1985.

[33] D.J. Gross, A. Neveu, J. Scherk, and J.H. Schwartz. Renormalization and unitarity in dual resonance models. *Phys. Rev. D*, 2:697–710, 1970.

[34] Murat Günaydin. Ads/cft dualities and the unitary representations of non-compact groups and supergroups: Wigner versus Dirac. In *Proc. International Wigner Symposium*, Istanbul, August 1999.

[35] R.W. Johnson. Finite integral equations for green's functions for : ϕ^4 : coupling. I. *J. Math. Phys.*, 11:2161, 1970.

[36] S. Kehrein, F. Wegner, and Yu.M. Pis'mak. Conformal symmetry and the spectrum of anomalous dimensions in the n-vector model in four - epsilon dimensions. *Nucl.Phys. B*, 402:669–692, 1993.

[37] O. Lauscher and M. Reuter. Ultraviolet fixed point and generalized flow equation of quantum gravity. *Phys. Rev. D*, 65:025013, 2002.

[38] J.D. Lawson. Polar and Ol'shanskii decompositions. *J. Reine Angewandte Mathematik*, 448:191–219, 1994.

[39] M. Lüscher. Operator product expansions on the vacuum in conformal quantum field theory in two space time dimensions. *Commun. Math. Phys.*, 50:23–52, 1976.

[40] M. Lüscher. *Theorie holomorpher Liehalbgruppen und deren Anwendung in konform kovarianten quantenfeldtheorien*. PhD thesis, Universität Hamburg, Germany, 1976.

[41] G. Mack. Conformal invariance and short distance expansions in quantum field theory. In *Strong interaction physics*, volume 17 of *Lecture Notes in Physics*, page 300, Heidelberg, 1972. Springer Verlag.

[42] G. Mack. Group theoretical approach to conformal invariant quantum field theory. *Journal de physique*, 34(C1 (Suppl. No. 10)):99, 1973.

[43] G. Mack. Group theoretical approach to conformal invariant quantum field theory. In E.R. Caianello, editor, *Renormalization and invariance in quantum field theory*, New York, 1974. Plenum Press.

[44] G. Mack. Osterwalder Schrader positivity in conformal invariant quantum field theory. In H. Rollnik and K. Dietz, editors, *Trends in elementary particle theory*, volume 37 of *Lecture Notes in Physics*, page 66. Springer Verlag, 1975.

[45] G. Mack. All unitary ray representations of the conformal group with positive energy. *Commun. Math. Phys.*, 55:1, 1977.

[46] G. Mack. Convergence of operator product expansions on the vacuum in conformal invariant quantum field theory. *Commun. Math. Phys.*, 53:155, 1977.

[47] G. Mack. Duality in quantum field theory. *Nuclear Physics*, B118:445, 1977.

[48] G. Mack and M. de Riese. Simple space time symmetries: generalizing conformal field theory. *J. Math. Phys.*, 48:052304, 2007.

[49] G. Mack and Martin Lüscher. Global conformal invariance. *Commun. Math. Phys.*, 41:203, 1975.

[50] G. Mack and T. Prüstel. Generalized gauge theories with nonunitary parallel transporters: General relativity with a cosmological constant as an example. *Eur. Phys. J. C*, 46:255–267, 2006.

[51] G. Mack and K. Symanzik. Currents, stress energy tensor and generalized unitarity in conformal invariant quantum field theory. *Commun. Math. Phys.*, 27:247, 1972.

[52] G. Mack and I.T. Todorov. Conformal invariant Green functions without ultraviolet divergences. *Physical Review*, D8:1764, 1971.

[53] A.A. Migdal. Conformal invariance and bootstrap. *Phys. Letters*, 37B:386, 1971.

[54] A.A. Migdal. On hadronic interactions at small distances. *Phys. Letters*, 37B:98, 1971.

[55] N.M. Nikolov, K.H. Rehren, and I. Todorov. Harmonic bilocal fields generated by globally conformal invariant scalar fields. *Commun. Math. Phys.*, 279:225–250, 2008.

[56] N.M. Nikolov, Y.S. Stanley, and I.T. Todorov. Globaly conformal invariant gauge field theory and rational correlation functions. *Nucl. Phys. B*, 670:373–400, 2003.

[57] K. Osterwalder and R Schrader. Axioms for euclidean green's functions, i. *Commun. Math. Phys.*, 31:83–112, 1973.

[58] K. Osterwalder and R Schrader. Axioms for euclidean green's functions, ii. *Commun. Math. Phys.*, 42:281–305, 1975.

[59] N.Ya. Palchik and E.S. Fradkin. A family of secondary fields in the d-dimensional conformal quantumk field theory. *Dokl. Phys.*, 44:757–759, 1999.

[60] N.Ya. Palchik and V.N. Zaikov. Secondary field in $d \geq 3$ conformal field theory: The existence condition for d-dimensional generalization of two-dimensional models in quantum field theory. In *Quantization, gauge theory and strings*, pages 375–381, Moscow, 2000.

[61] G. Parisi and L. Peliti. Calculation of critical indices. *Lettere Nuovo Cimento*, 2:386, 1971.

[62] Yu. M. Pis'mak. Renormalization group and infinite algebraic structure in D-dimensional conformal field theory. *J. Phys.*, A39:8157–8172, 2006.

[63] K.H. Rehren. Algebraic holography. *Annales Henri Poincare*, 1:607–623, 2000.

[64] W Rühl. *The Lorentz group and harmonic analysis*. Benjamin Inc., 1970.

[65] W. Rühl. Lifting a conformal field theory from d-dimensional flat space to $d+1$ dimensional ads-space. *Nucl.Phys. B*, 705:437–456, 2005.

[66] E.R. Speer. *Generalized Feynman amplitudes*. Princeton University Press, 1969.

[67] R.F. Streater and A.S. Wightman. *PCT, Spin and Statistics, and All That*. Addison-Wesley Publishing Company, 1964, 1980.

[68] L. Susskind and J. Lindesay. *An introduction to black holes, information and the string theory revolution: The holographic universe*. World Scientific, 2005.

[69] K. Symanzik. Green's functions method and renormalization of renormalizable quantum field theories. In B. Jaksic, editor, *Lectures on High Energy Physics*, pages 485–517, New York, 1966. Gordon and Breach. Proceedings Sixth Summer Meeting of Nuclear Physicists, Hercegnovi 1961.

[70] K. Symanzik. On calculations in conformal invariant field theories. *Nuovo Cimento*, 3:734–738, 1972.

[71] A.N. Vasiliev, Yu.M. Pis'mak, and Yu.R. Khonkonen. 1/n expansion. calculation of the exponent eta in the order 1/n**3 by the conformal bootstrap. *Theor. Math. Phys.*, pages 127–134, 1982.

[72] N.N. Vasiliev. *Quantum field renormalization group in the theory of critical behavior and stochastic dynamics*. Chapmann and Hall, London, 2004.

[73] G. Veneziano. Construction of crossing-symmetric, Regge behaved amplitude for linearly rising trajectories. *Nuovo Cimento*, 57:190, 1968.

[74] G. Veneziano. Narrow-resonance models compatible with duality and their developments. In A. Zichichi, editor, *Elementary processes at high energy*, pages 94–158. Academic Press, 1971. Proc. Erice summer school 1970.

[75] K. Wilson. Non-Lagrangean models of current algebra. *Phys.Rev.*, 179:1499, 1969.

[76] K. Wilson. Operator product expansion and anomalous dimensions in the Thirring model. *Phys.Rev. D*, 2:1473, 1970.

[77] K. G. Wilson and John B. Kogut. The Renormalization group and the epsilon expansion. *Phys. Rept.*, 12:75–200, 1974.

[78] E. Witten. (2+1)-dimensional gravity as a exactly soluble system. *Nucl. Phys. B*, 311:46, 1988.